

HIAS-E-107

**A new test for common breaks
in heterogeneous panel data models**

Peiyun Jiang^(a), Eiji Kurozumi^(b)

(a) *Hitotsubashi Institute for Advanced Study, Hitotsubashi University*

(b) *Department of Economics, Hitotsubashi University*

May 2021



Hitotsubashi Institute for Advanced Study, Hitotsubashi University
2-1, Naka, Kunitachi, Tokyo 186-8601, Japan
tel:+81 42 580 8668 <http://hias.hit-u.ac.jp/>

HIAS discussion papers can be downloaded without charge from:
<https://hdl.handle.net/10086/27202>
<https://ideas.repec.org/s/hit/hiasdp.html>

All rights reserved.

A new test for common breaks in heterogeneous panel data models ¹

Peiyun Jiang

Eiji Kurozumi

Hitotsubashi Institute for Advanced Research

Department of Economics

Hitotsubashi University

Hitotsubashi University

This version: May 20, 2021

Abstract

In this paper, we develop a new test to detect whether break points are common in heterogeneous panel data models where the time series dimension T could be large relative to cross-section dimension N . The error process is assumed to be cross-sectionally independent. The test is based on the cumulative sum (CUSUM) of ordinary least squares (OLS) residuals. We derive the asymptotic distribution of the detecting statistic under the null hypothesis, while proving the consistency of the test under the alternative. Monte Carlo simulations and an empirical example show good performance of the test.

JEL classification: C12, C23

Key words: CUSUM test, panel data, structural change, common breaks

¹Kurozumi's research was supported by JSPS KAKENHI Grant Number 19K01585. Address correspondence to Peiyun Jiang, Hitotsubashi Institute for Advanced Research, Hitotsubashi University, 2-1 Naka, Kunitachi, Tokyo 186-8601, Japan; e-mail: pynjiang@gmail.com

1. Introduction

In recent years, panel data models have become increasingly popular in theoretical and empirical analyses, since richer information from both the cross-section and time series dimension leads to more powerful inferences than with a single cross-section or a single time series. In particular, the modeling and inferences of structural changes in panel frameworks have attracted significant attention in the literature. Compared to applying the single detection method for structural changes separately to each series, using cross-sectional datasets improves break detection power. The detection procedures in panels are often designed to test for the null hypothesis that the regression parameters in each series are constant over time against the alternative that at least one series exhibits structural changes. See, for example, Horváth and Hušková (2012) in a mean-shift panel model, De Wachter and Tzavalis (2012) and Hidalgo and Schafgans (2017) in dynamic panels, Pauwels *et al.* (2012) in panel data models allowing for heterogeneous coefficients, Chen and Huang (2018) in a time-varying panel data model, and Antoch *et al.* (2019) in panels with fixed T and large N, to name a few. However, the rejection of the null hypothesis leaves the researcher with no information as to which cross-sectional unit exhibits structural changes. Furthermore, it naturally leads to the issue of change point estimation in panel data models.

Classical change point estimation methodologies in panel literature often assume that break point occurred in each series at the same location, referred to as the common break point. This assumption is particularly attractive, as the common break phenomenon occurs in many practical applications. The other major advantage of this assumption is the increased accuracy of the change point estimate, as noted by Bai (2010). It is well known that only the break fraction (i.e., the break date divided by the sample size) can be consistently estimated in a single time series. In panel frameworks, however, the failure of the consistency of the break point in time series models has been overcome under the common break assumption. This enhanced precision of the common break point estimate has been widely confirmed under various frameworks in panel data analyses. Kim (2011, 2014) focused on panel deterministic time trend models and considered a factor structure for the error component. Although the former study stated that the ordinary least squares break date estimator fails to achieve

consistency by imposing the factor structure, the latter overcame this problem and developed a new estimation strategy, where the common break date is estimated jointly with the common factor to successfully sustain the precision advantage of the common break point estimate in panels. In addition, Qian and Su (2016) used a panel data model in which the parameters of interest are homogeneous and errors are assumed to be cross-sectionally independent, while Baltagi *et al.* (2016) considered a more general panel framework allowing for heterogeneous parameters across individuals and multifactor error structure. More related works, including Li *et al.* (2016), Baltagi *et al.* (2017), Horváth *et al.* (2017), Westerlund (2019), and others, have documented that the break date estimate obtains increased precision via imposing a common break assumption in panels.

In practice, however, the common break assumption is restrictive, and some evidence has verified that the break points are likely to vary significantly across individuals (see Claeys and Vašíček 2014; Adesanya 2020). To the best of our knowledge, no study has focused on the validity of the common break assumption in panels. In this paper, we contribute to the literature in three ways. First, we fill in this gap to introduce a test for the null hypothesis that the panels exhibit a common break against the alternative that break dates can vary across units. The closest related work is that of Oka and Perron (2018), who considered common break detection in maximum likelihood frameworks in multiple equation systems. We extend their model to a more general framework where both the number of series N and the number of observations T are sufficiently large, which makes it available using panel or macroeconomic data in applications.

The second major contribution of this paper is that we investigate the statistical properties of the estimated common break point when the common break assumption fails. It is verified that the common break estimate cannot be consistent for each series, but will be restricted to a specific region. Based on this property, our test has a non-degenerate distribution under the null hypothesis and achieves consistency under the alternative.

Third, our test delivers monotonic power as the magnitude of the breaks increases. The statistic is established by the squares of the cumulative sum of the residuals, and we use a normalization factor to replace the long-run variance estimator to avoid power loss when

the shift increases under the alternative (the so-called nonmonotonic power problem). Monte Carlo simulations show good size performance for large T . Moreover, the test can successfully reject the null hypothesis of a common break against various types of alternatives and has nontrivial power for large breaks. An empirical example demonstrates that a common break exists in the mutual fund data during the sub-prime crisis.

From a different perspective, recent clustering literature suggested an estimation methodology as an alternative strategy to identify distinct breaks across units in panels. The panel data are modeled using a grouped pattern, in which the regression coefficients containing break dates are heterogeneous across groups but homogeneous within a group. In this framework, Okui and Wang (2020) and Lumsdaine *et al.* (2020) proposed iterative estimation approaches to jointly estimate the break point, group membership structure, and coefficients. The consistency of all estimates can be achieved simultaneously within the prior information on the number of groups and an appropriate choice of the initial values for iteration. Researchers can determine whether to conduct a testing procedure, apply an estimation methodology, or use a hybrid of two approaches depending on their empirical purpose.

The remainder of this paper is organized as follows. Section 2 introduces the model and necessary assumptions. Section 3 explains the testing strategy for the common break assumption. Section 4 establishes the asymptotic distribution of the statistic under the null hypothesis and the consistency of the test under the alternative hypothesis. Monte Carlo simulations are conducted in Section 5. Section 6 provides an empirical example, and Section 7 provides concluding remarks. The mathematical proofs are relegated to the Appendix.

2. Model and Assumptions

We consider a panel data model allowing for heterogeneous coefficients across units, defined by

$$y_{it} = x'_{it}\beta_i + x'_{it}\delta_i 1_{\{t > k_i^0\}} + u_{it}, \quad 1 \leq i \leq N \text{ and } 1 \leq t \leq T, \quad (1)$$

where $x_{it} = [x_{it}(1), \dots, x_{it}(p)]'$ is p -dimensional explanatory variables including a constant term; thus, the first element is unity for all t . The coefficients $\beta_i = [\beta_{i1}, \dots, \beta_{ip}]'$, $\delta_i = [\delta_{i1}, \dots, \delta_{ip}]'$ are $p \times 1$ vectors of fixed parameters, and $1_{\{t > k_i^0\}}$ is an indicator function that

takes the value one if $t > k_i^0$, and zero otherwise. u_{it} is an unobservable stochastic disturbance. We assume that the regression parameters in the i th panel change from β_i to $\beta_i + \delta_i$ at unknown time k_i^0 , and we are interested in testing whether the break point in each series is common against the alternative that the break point varies across individuals. The null hypothesis is defined as

$$H_0 : k_i^0 = k^0, \quad \text{for all } i = 1, 2, \dots, N.$$

Under the alternative of distinct breaks across individuals, we suppose that there exist G groups, and the regression coefficients share the common break point in each group $g = 1, 2, \dots, G$. Then, the alternative hypothesis is defined by

$$H_A : k_{g_1}^0 \neq k_{g_2}^0, \quad \text{for some } g_1, g_2 \in \{1, 2, \dots, G\}.$$

In this paper, we impose the following assumptions.

Assumption 1 $k_i^0 = [T\tau_i^0]$, where $\tau_i^0 \in (0, 1)$ and $[\cdot]$ is the greatest integer function.

The break point k_i^0 , which is a positive fraction of the total sample size, is assumed to be bounded away from the end points. This is a conventional assumption in the change point literature, see Bai (1997).

Assumption 2 Define $\phi_N = \sum_{i=1}^N \delta_i^{0'} \delta_i^0$. Suppose that

- (i) $\phi_N \rightarrow \infty$ as $N \rightarrow \infty$,
- (ii) $\frac{\phi_N}{N}$ is bounded as $N \rightarrow \infty$,
- (iii) $\frac{T}{N} \rightarrow \infty$, $\phi_N \frac{\sqrt{T}}{N} \rightarrow \infty$ as $(T, N) \rightarrow \infty$.

Denote δ_i^0 as the true shift for individual i . Assumptions 2(i)–(ii) are borrowed from Assumption A2 in Baltagi *et al.* (2016). The additional condition $T/N \rightarrow \infty$ requires that T grows at a faster rate than N . This is a significant condition to ensure a non-degenerate distribution of the statistic under the null hypothesis and consistency of the test under the alternative.

Assumption 3 (i) For each series i , u_{it} is independent of x_{it} for all i and t ;

(ii) $u_{it} = \sum_{j=0}^{\infty} a_{ij} \epsilon_{i,t-j}$, $\epsilon_{it} \sim (0, \sigma_{i\epsilon}^2)$ are *i.i.d* over all i and t ; $\sum_j j |a_{ij}| \leq M$ for all i .

The idiosyncratic errors form a stationary time series, and it is assumed that u_{it} are cross-sectionally independent, similar to the assumption in Bai (2010). In practice, this assumption is relatively restrictive, as cross-sectional dependence commonly exists in many panel datasets. As explained in Sections 3 and 4, the statistic of our test can have a non-degenerate distribution under the null hypothesis of the common break, crucially depending on the consistency of the common change point estimate. However, Kim (2011) indicated that imposing a factor structure on the error component may impede the consistency property. Some additional techniques are needed if we relax Assumption 3 to allow for cross-sectional dependence.

Assumption 4 (i) For $i = 1, \dots, N$, the matrices $(1/j) \sum_{t=1}^j x_{it}x'_{it}$, $(1/j) \sum_{t=T-j+1}^T x_{it}x'_{it}$, $(1/j) \sum_{t=k_i^0-j+1}^{k_i^0} x_{it}x'_{it}$, and $(1/j) \sum_{t=k_i^0+1}^{k_i^0+j} x_{it}x'_{it}$ are stochastically bounded and have minimum eigenvalues uniformly bounded away from zero in probability for all large j .

(ii) For each i , $(1/T) \sum_{t=1}^T x_{it}x'_{it}$ converges in probability to a nonrandom and positive definite $p \times p$ matrix C_i as $T \rightarrow \infty$.

(iii) For each i , $(1/T) \sum_{t=1}^T x_{it}$ converges in probability to a $p \times 1$ vector c_{i1} as $T \rightarrow \infty$.

Denote the j th row of C_i by c_{ij} for $j = 1, \dots, p$. That is, $C = [c_{i1}, \dots, c_{ip}]'$. Note that the vector c'_{i1} is the first row of C_i .

Assumption 5 (i) For any positive finite integer s , the matrices $(1/N) \sum_{i=1}^N \sum_{t=k_i^0-s+1}^{k_i^0} x_{it}x'_{it}$ and $(1/N) \sum_{i=1}^N \sum_{t=k_i^0+1}^{k_i^0+s} x_{it}x'_{it}$ are stochastically bounded and have minimum eigenvalues uniformly bounded away from zero in probability for all large N .

(ii) For each t , $(1/N) \sum_{i=1}^N x_{it}x'_{it}$ is stochastically bounded as $N \rightarrow \infty$.

Assumption 4 is a conventional assumption in time series models, see, for example, Bai (1997), while Assumption 5 is borrowed from Assumption 5 in Baltagi *et al.* (2016).

3. Test Statistic

The null hypothesis assumes that the panels exhibit one break occurring at an unknown common location. We first use the least squares method, as proposed by Baltagi *et al.* (2016),

to estimate the common break point. Let

$$Y_i = \begin{bmatrix} y_{i1} \\ y_{i2} \\ \vdots \\ y_{iT} \end{bmatrix}, X_i = \begin{bmatrix} x'_{i1} \\ x'_{i2} \\ \vdots \\ x'_{iT} \end{bmatrix}, Z_i(k_i) = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ x'_{i(k_i+1)} \\ \vdots \\ x'_{iT} \end{bmatrix}, \text{ and } u_i = \begin{bmatrix} u_{i1} \\ u_{i2} \\ \vdots \\ u_{iT} \end{bmatrix}.$$

The model with an unknown break point k_i can be rewritten in matrix form as

$$\begin{aligned} Y_i &= X_i\beta_i + Z_i(k_i)\delta_i + u_i \\ &= [X_i, Z_i(k_i)] \begin{bmatrix} \beta_i \\ \delta_i \end{bmatrix} + u_i \\ &= \bar{X}_i(k_i)b_i + u_i, \end{aligned} \tag{2}$$

where $\bar{X}_i(k_i) = [X_i, Z_i(k_i)]$, and $b_i = [\beta_i', \delta_i']'$. Given any $k^* = 1, 2, \dots, T-1$, b_i can be estimated by

$$\hat{b}_i(k^*) = \begin{bmatrix} \hat{\beta}_i(k^*) \\ \hat{\delta}_i(k^*) \end{bmatrix} = [\bar{X}_i(k^*)' \bar{X}_i(k^*)]^{-1} \bar{X}_i(k^*)' Y_i, \quad i = 1, \dots, N.$$

The sum of squared residuals for i th equation is given by

$$SSR_i(k^*) = [Y_i - \bar{X}_i(k^*)\hat{b}_i(k^*)]' [Y_i - \bar{X}_i(k^*)\hat{b}_i(k^*)], \quad i = 1, \dots, N.$$

The least squares estimator of k^* is defined as

$$\hat{k} = \arg \min_{1 \leq k^* \leq T-1} \sum_{i=1}^N \pi_i SSR_i(k^*). \tag{3}$$

where weights $\pi_i \in (0, 1)$, $i = 1, \dots, N$, $\sum_{i=1}^N \pi_i = 1$.

Our statistic is composed of ordinary least squares residuals based on the estimated common break point \hat{k} . We decompose the panels into two regimes using \hat{k} in the time series dimension. Then, the OLS residuals are calculated by

$$\hat{u}_i = \begin{bmatrix} \hat{u}_{i1} \\ \hat{u}_{i2} \\ \vdots \\ \hat{u}_{iT} \end{bmatrix} = Y_i - \bar{X}_i(\hat{k})\hat{b}_i(\hat{k}), \tag{4}$$

and the squares of the partial sum of the OLS residuals \hat{u}_{it} are defined by

$$US_{NT}(k, \hat{k}) = \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k \hat{u}_{it} \right)^2, \quad \text{where } k = [T\tau] \text{ with } \tau \in (0, 1). \quad (5)$$

The statistic is a CUSUM-type of residuals, motivated by the consistency of the break point estimate if the common break assumption holds. Under the null hypothesis that all individuals are assumed to share a common change point $k^0 = [T\tau^0]$ with $\tau^0 \in (0, 1)$, Baltagi *et al.* (2016) verified that the common break date is consistently estimated. Based on the consistency of $\hat{k} \xrightarrow{p} k^0$, the regression parameters corresponding to regimes $\{x_{i1}, \dots, x_{i\hat{k}}\}$, $\{x_{i(\hat{k}+1)}, \dots, x_{iT}\}$ are asymptotically constant over time. Consequently, the cumulative sums of the corresponding residuals will not diverge and can have a non-degenerate distribution, which is derived as follows:

$$US_{NT}(k, \hat{k}) \Rightarrow \begin{cases} \sigma^2 [W(\tau) - \frac{\tau}{\tau^0} W(\tau^0)]^2 & \text{if } \tau \leq \tau^0 \\ \sigma^2 [W(\tau) - W(\tau^0) - \frac{\tau - \tau^0}{1 - \tau^0} (W(1) - W(\tau^0))]^2 & \text{if } \tau > \tau^0 \end{cases},$$

where $W(\cdot)$ is a one-dimensional Brownian motion, and σ^2 is the long-run variance defined below. Under the alternative of distinct breaks, since the estimated common break point cannot coincide with the true break point for each series, partial residuals will significantly deviate from the one under the null hypothesis. Hence, $US_{NT}(k, \hat{k})$ will diverge to infinity as $N, T \rightarrow \infty$ such that we can successfully reject the null hypothesis.

A traditional approach is to use a consistent estimate to replace the unknown σ^2 , while the kernel estimator is commonly applied. Typically, the selection of the bandwidth for the kernel estimator significantly affects the size and power performance of the test. In time series analyses, it has been extensively mentioned that the structural change tests suffer from the so-called non-monotonic power problem; that is, the tests may lose power as the magnitude of the break increases. See Vogelsang (1999), Deng and Perron (2008), Yamazaki and Kurozumi (2015), and Jiang and Kurozumi (2019), among others. The main reason is that the long-run variance estimated under the null hypothesis is consistent but may be severely biased under the alternative hypothesis. To maintain nontrivial detection power for large breaks, we extend the self-normalization method proposed by Shao and Zhang (2010) to construct a normalization factor instead of using the long-run variance estimate. This normalization

factor $V_{NT}(k_1, \hat{k}, k_2)$ is required to be proportional to σ^2 such that the long-run variance can be canceled out as

$$\frac{US_{NT}(k, \hat{k})}{V_{NT}(k_1, \hat{k}, k_2)} \Rightarrow \frac{\sigma^2 \text{ functional of Brownian motions}}{\sigma^2 \text{ functional of Brownian motions}},$$

where the long-run variance σ^2 is $\lim_{(N,T) \rightarrow \infty} E \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T u_{it} \right)^2$. Furthermore, the normalization process cannot grow at a faster rate relative to the process $US_{NT}(k, \hat{k})$ under the alternative to avoid loss of power. To this end, we separate the panels into four regimes by flexible points k_1, k_2 and the estimated break point \hat{k} , where k_1 and k_2 take values in the interval $1 \leq k_1 < \hat{k} < k_2 \leq T - 1$. We estimate the model on the basis of four regimes $\{x_{i1}, \dots, x_{ik_1}\}$, $\{x_{i(k_1+1)}, \dots, x_{i\hat{k}}\}$, $\{x_{i(\hat{k}+1)}, \dots, x_{ik_2}\}$, and $\{x_{i(k_2+1)}, \dots, x_{iT}\}$ for the i th equation. Denote $T \times p$ matrices by

$$X_{ji}(a, b) = [0, \dots, 0, x_{i,a+1}, \dots, x_{i,b}, 0, \dots, 0]', \quad j = 1, 2, \quad (6)$$

$$X_{3i}(a) = [0, \dots, 0, x_{i,a+1}, \dots, x_{iT}]', \quad (7)$$

where the elements of the $(a + 1)$ th- b th rows of $X_{ji}(a, b)$ are the same as that of X_i and zero otherwise, and the elements of the $(a + 1)$ th- T th rows of $X_{3i}(a)$ are the same as that of X_i and zero otherwise. Then, the model can be represented by

$$\begin{aligned} Y_i &= [X_i, X_{1i}(k_1, \hat{k}), X_{2i}(\hat{k}, k_2), X_{3i}(k_2)] \begin{bmatrix} \beta_i \\ \delta_{1i} \\ \delta_{2i} \\ \delta_{3i} \end{bmatrix} + u_i \\ &= X_i \beta_i + X_{1i}(k_1, \hat{k}) \delta_{1i} + X_{2i}(\hat{k}, k_2) \delta_{2i} + X_{3i}(k_2) \delta_{3i} + u_i \\ &= \tilde{X}_i(k_1, \hat{k}, k_2) b_i + u_i, \end{aligned} \quad (8)$$

where $\tilde{X}_i(k_1, \hat{k}, k_2) = [X_i, X_{1i}(k_1, \hat{k}), X_{2i}(\hat{k}, k_2), X_{3i}(k_2)]$. Using the coefficient estimators $\tilde{\beta}_i$, $\tilde{\delta}_{1i}$, $\tilde{\delta}_{2i}$, and $\tilde{\delta}_{3i}$, the corresponding residuals are calculated as

$$\tilde{u}_i = \begin{bmatrix} \tilde{u}_{i1} \\ \tilde{u}_{i2} \\ \vdots \\ \tilde{u}_{iT} \end{bmatrix} = Y_i - X_i \tilde{\beta}_i - X_{1i}(k_1, \hat{k}) \tilde{\delta}_{1i} - X_{2i}(\hat{k}, k_2) \tilde{\delta}_{2i} - X_{3i}(k_2) \tilde{\delta}_{3i}. \quad (9)$$

Then, we define the process $V_{NT}(k_1, \hat{k}, k_2)$ based on the residuals \tilde{u}_{it} as

$$\begin{aligned}
& V_{NT}(k_1, \hat{k}, k_2) \\
&= \frac{1}{T} \sum_{s=1}^{k_1} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^s \tilde{u}_{it} \right)^2 + \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} \tilde{u}_{it} \right)^2 \\
&+ \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s \tilde{u}_{it} \right)^2 + \frac{1}{T} \sum_{s=k_2+1}^T \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T \tilde{u}_{it} \right)^2. \tag{10}
\end{aligned}$$

Thus, our test statistic is composed of the squared CUSUM of residuals (5) and the normalization factor (10), defined by

$$\begin{aligned}
& S_{NT}(k, k_1, k_2) = \sup_{(k, k_1, k_2) \in \Omega(\epsilon)} \frac{US_{NT}(k, \hat{k})}{V_{NT}(k_1, \hat{k}, k_2)} \\
&= \sup_{(k, k_1, k_2) \in \Omega(\epsilon)} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k \hat{u}_{it} \right)^2 \left\{ \frac{1}{T} \sum_{s=1}^{k_1} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^s \tilde{u}_{it} \right)^2 \right. \\
&+ \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} \tilde{u}_{it} \right)^2 + \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s \tilde{u}_{it} \right)^2 \\
&\left. + \frac{1}{T} \sum_{s=k_2+1}^T \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T \tilde{u}_{it} \right)^2 \right\}^{-1},
\end{aligned}$$

where $\Omega(\epsilon) = \{(k, k_1, k_2) \text{ or } (\tau, \tau_1, \tau_2) : [T\epsilon] \leq k \leq [T(1-\epsilon)], [T\epsilon] \leq k_1 \leq \hat{k} - [T\epsilon], \hat{k} + [T\epsilon] \leq k_2 \leq [T(1-\epsilon)]\}$. $k = [T\tau]$, $k_1 = [T\tau_1]$ and $k_2 = [T\tau_2]$ with $\tau, \tau_1, \tau_2 \in (0, 1)$.

4. Asymptotic Theory

We next derive the limiting properties of the test statistic.

Theorem 1 *Suppose that Assumptions 1–5 hold. Then, under H_0 , we have, as $N, T \rightarrow \infty$,*

$$\begin{aligned}
& S_{NT}(k, k_1, k_2) \\
\Rightarrow & \sup_{(\tau, \tau_1, \tau_2) \in \Omega(\epsilon)} \left\{ W(\tau) - \tau \frac{W(\tau^0)}{\tau^0} - (\tau - \tau^0) \left[\frac{W(1) - W(\tau^0)}{1 - \tau^0} - \frac{W(\tau^0)}{\tau^0} \right] 1_{\{\tau > \tau^0\}} \right\}^2 \\
& \left\{ \int_0^{\tau_1} \left[W(s) - s \frac{W(\tau_1)}{\tau_1} \right]^2 ds + \int_{\tau_1}^{\tau_0} \left[W(\tau^0) - W(s) - (\tau^0 - s) \frac{W(\tau^0) - W(\tau_1)}{\tau^0 - \tau_1} \right]^2 ds \right. \\
& + \int_{\tau^0}^{\tau_2} \left[W(s) - W(\tau^0) - (s - \tau^0) \frac{W(\tau_2) - W(\tau^0)}{\tau_2 - \tau^0} \right]^2 ds \\
& \left. + \int_{\tau_2}^1 \left[W(1) - W(s) - (1 - s) \frac{W(1) - W(\tau_2)}{1 - \tau_2} \right]^2 ds \right\}^{-1},
\end{aligned}$$

where $W(\cdot)$ is a standard Brownian motion, $k = [T\tau]$, $k_1 = [T\tau_1]$, $k^0 = [T\tau^0]$, and $k_2 = [T\tau_2]$ with $\tau, \tau_0, \tau_1, \tau_2, \in (0, 1)$.

Under the null hypothesis, the proposed test has a non-standard limit distribution depending on the true break fraction, which is unknown in practice. We choose $\tau^0 = 0.1, 0.2, \dots, 0.9$, and approximate Brownian motions using 2,000 independent normal random variables with 10,000 replications to obtain the critical values in Table 1. A researcher can calculate an appropriate critical value based on the value of the estimated break fraction. For example, if $\hat{\tau} \in [0.4, 0.5)$, we obtain the critical value by the interpolation,

$$c = c_{0.4} + 10(\hat{\tau} - 0.4)(c_{0.5} - c_{0.4}),$$

where c_{τ^0} for $\tau^0 = 0.1, \dots, 0.9$ are the critical values given in Table 1. Next, we investigate the behavior of the proposed test statistic when the breaks vary across individuals. We focus on the case in which there are two groups, and individuals in the same group share a common break $k_j^0, j = 1, 2$.

$$H_{1A} : |k_1^0 - k_2^0| \geq \Delta T, \text{ for some } \Delta > 0.$$

Assumption 6 *Let $N_j, j = 1, 2$, denote the number of units in group j ($N = N_1 + N_2$). Suppose that $N_j/N \rightarrow \pi_j > 0$ for $j = 1, 2$.*

To characterize the limiting properties of the test statistic under the alternative, it is useful to first state some preliminary results regarding the statistical properties of the estimated

common break point. Define $K(C) = \{k : 1 \leq k < k_1^0 - C_1, k_2^0 + C_2 < k \leq T - 1\}$, where C_1, C_2 are finite numbers.

Proposition 1 *Suppose that Assumptions 1–6 hold. Then, under H_{1A} , for any given $\epsilon > 0$, for both large N and T ,*

$$P(\hat{k} \in K(C)) < \epsilon.$$

Proposition 1 states the possible region of the location of the common break date estimator when the common break assumption fails. This implies that this estimator will be bounded away from both end points. In other words, the estimated common break point may lie between the two true break points or be stochastically bounded by either of the true break dates.

Proposition 2 *Suppose that Assumptions 1–6 hold. Under H_{1A} , for both large N and T ,*

(i) *if $\hat{k} < k_1^0$,*

$$\sup_{k \in \Omega(\epsilon)} US_{NT}(k, \hat{k}) = O_p(NT),$$

(ii) *if $k_1^0 \leq \hat{k} \leq k_2^0$,*

$$\sup_{k \in \Omega(\epsilon)} US_{NT}(k, \hat{k}) = O_p(NT),$$

(iii) *if $k_2^0 < \hat{k}$,*

$$\sup_{k \in \Omega(\epsilon)} US_{NT}(k, \hat{k}) = O_p(NT),$$

Proposition 2 derives the divergence rate of the process $US_{NT}(k, \hat{k})$ under the alternative. In case (i), Proposition 1 implies that the common change point estimate is bounded by the true break point k_1^0 ; that is, $k_1^0 - \hat{k} = O_p(1)$. Since we assume that the two true breaks are separated by some positive fraction of the sample size, \hat{k} will become distant from the other break date k_2^0 . Therefore, for individuals in group 2, the regression parameters will be estimated based on an inconsistent break fraction estimate. Then, we find that the CUSUM of the corresponding residuals \hat{u}_{it} in $US_{NT}(k, \hat{k})$ will diverge to infinity at a rate of NT . For the second and third cases, it is shown that the divergence rate of the process $US_{NT}(k, \hat{k})$ is the same as that in case (i).

Proposition 3 *Suppose that Assumptions 1–6 hold. Under H_{1A} , for any given $\epsilon > 0$, there exists a finite $M > 0$ such that, for both large N and T ,*

(i)

$$P\left(\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) > M \mid k_1^0 - C_1 < \hat{k} \leq k_1^0\right) < \epsilon,$$

(ii)

$$P\left(\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) > M \mid k_1^0 < \hat{k} < k_2^0\right) < \epsilon,$$

(iii)

$$P\left(\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) > M \mid k_2^0 \leq \hat{k} < k_2^0 + C_2\right) < \epsilon.$$

Proposition 3 investigates the limiting properties of the normalization process under the alternative. The results indicate that $\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2)$ is $O_p(1)$. Since the model is estimated based on four subsamples for the normalization factor, we can eventually find appropriate k_1 and k_2 such that the minimization will not diverge. The numerator of the statistic diverges at a rate of NT , and the denominator has a finite limit. Then, we derive the consistency of the test under the alternative in the following theorem:

Theorem 2 *Suppose that Assumptions 1–6 hold. Then, under H_{1A} , we have, as $N, T \rightarrow \infty$,*

$$S_{NT}(k, k_1, k_2) \rightarrow \infty.$$

The consistency of this test is achieved under a particular and specified alternative H_{1A} . Nevertheless, our simulations confirm that this test is valid and powerful against a variety of alternatives.

5. Finite Sample Properties

In this section, we investigate the finite sample performance of the test considered in the previous sections. The data-generating process (DGP.1) under the null hypothesis of a common break is given by

$$y_{it} = x'_{it}\beta_i + x'_{it}\delta_i 1_{\{t > k^0\}} + u_{it}, \quad i = 1, \dots, N, \quad t = 1, \dots, T.$$

where $x_{it} = [1, z_{it}]'$ includes a constant, each z_{it} has a normal distribution $N(1, 1)$, and is independent of the errors u_{it} , $1 \leq t \leq T, 1 \leq i \leq N$. We assume that a common break $k^0 = [0.5T]$ exists in the slopes. The coefficients $\beta_i \sim i.i.d.U(0, 0.8)$ and δ_i are the jumps for each series with $\delta_i \sim i.i.d.U(0, 0.5)$. We allow for serial correlation in the errors $u_{it} = \rho u_{i(t-1)} + e_{it}$ with $e_{it} \sim i.i.d.N(0, (1 - \rho)^2)$. The trimming parameter ϵ is 0.1, the number of replications is 2,000, and all computations are conducted using the GAUSS matrix language.

Table 2 summarizes the empirical sizes of the test for the different pairs of (N, T) . In the case of *i.i.d.* errors, the nominal rejection rate is close to the corresponding significance level of the test. When the errors are allowed to be serially correlated with $\rho = 0.4$, for small N and T , the size distortion is quite noticeable. The size improves for large T and appears to be quite close to the nominal level at $T = 200$.²

In practice, no prior information is available on the form of structural changes for researchers. Therefore, we conduct extensive simulations to explore the empirical power of the test for various group patterns of structural change and different magnitudes of the break. We first impose a benchmark case as Assumption 6. There are two groups in panels, and the break points for individuals are common in the same group but distinct across groups. We next consider more general circumstances in which there are more than two groups in panels or the break dates can be distinct across individuals. Moreover, we are interested in the validity of the test when the break dates are common, but multiple common breaks occurred in the panels. Then, four types of alternative hypotheses are considered as follows:

- H_{1A} : There are two groups and the series in each group share common break k_j^0 , $j = 1, 2$. Let N_j denote the number of units in group j and $N = N_1 + N_2$.
- H_{2A} : There are three groups and the series in each group share common break k_j^0 , $j = 1, 2, 3$. Note that $N = N_1 + N_2 + N_3$.
- H_{3A} : Suppose that there is no group pattern. The break point for the j th series is given by k_j^0 , $j = 1, 2, \dots, N$.

²The size is distorted corresponding to strong serial correlation ($\rho = 0.8$) but appears to be controlled when T increases. The results are similar and are thus omitted.

- H_{4A} : The panel data exhibit multiple common break dates.

The data generating process (DGP.2) under H_{1A} is given by

$$\begin{cases} y_{it} = x'_{it}\beta_i + x'_{it}\delta_{1i}1_{\{t>k_1^0\}} + u_{it} & t = 1, \dots, T, \text{ for } i \text{ in group 1,} \\ y_{it} = x'_{it}\beta_i + x'_{it}\delta_{2i}1_{\{t>k_2^0\}} + u_{it} & t = 1, \dots, T, \text{ for } i \text{ in group 2,} \end{cases}$$

which is the same as DGP.1, except that the change point varies across groups. The first group exhibits one common break at $k_1^0 = [0.25T]$, and we set the time of change k_2^0 equal to $[0.75T]$ in the second group. We assume $\beta_i \sim i.i.d.U(0, 0.8)$ and the jumps $\delta_{1i}, \delta_{2i} \sim i.i.d.U(0, 0.5)$. The ratio of units among the groups is set to $N_1 : N_2 = 5 : 5$. Table 3 shows that the test is powerful (almost with a rejection probability of more than 80%), except for small N or T. Table 4 reports the effect of the magnitude of change on power. As expected, the proposed test delivers monotonic power. When the magnitude of the changes is larger than 0.4, our test almost perfectly rejects the null hypothesis (power tends to one). We can see that the test shows good performance in the case of two well-separated groups. We further investigate the sensitivity of the test when the group characteristics (distance between two change points or number of units in each group) change. In Table 5, we fix one common break at $[0.2T]$, and the other break changes from $[0.25T]$ to $[0.8T]$. If the distance between two breaks exceeds $[0.3T]$, the rejection probability reaches at least 90% at the 10% significance level. When the two break dates become quite close (the distance is less than $[0.1T]$), the power of the test decreases to 0.325 at the 10% significance level. On the other hand, the power of the test is sensitive to the number of individuals in each group. Table 6 shows that the test rejects the null hypothesis with probability over 80% when the number of observations in each group is sufficiently large (the ratio of units between two groups is larger than 3/7). If the number of individuals in one group is much less than that in the second group, the heterogeneity between the two groups cannot be identified. Eventually, it is not easy to reject the null hypothesis, even if the two break dates are distinct.

We next investigate the power properties of the test under H_{2A} . The results for distinct change point locations and ratios of units among groups are reported in Table 7. The test can successfully reject the null of one common break for large N. The close break points among the three groups will reduce the power.

The alternative hypothesis H_{3A} considers heterogeneous change points without a group pattern. The change point for individual j is set to $k_j^0 = [T\tau_j^0]$, $j = 1, 2, \dots, N$, while the break fraction τ_j^0 is drawn from $U(0.15, 0.75)$. Table 8 shows that the test is still powerful for large N .

The data generating process (DGP.3) under H_{4A} is given by

$$y_{it} = x'_{it}\beta_i + x'_{it}\delta_{1i}1_{\{k_1^0 < t \leq k_2^0\}} + x'_{it}\delta_{2i}1_{\{t > k_2^0\}} + u_{it}, \quad t = 1, \dots, T, \quad i = 1, \dots, N,$$

where the coefficients change from β_i to $\beta_i + \delta_{1i}$ in the second regime and change from $\beta_i + \delta_{1i}$ to $\beta_i + \delta_{2i}$ in the third regime. The change points are set to $k_1^0 = [0.25T]$ and $k_2^0 = [0.75T]$, while the coefficients $\beta_i \sim i.i.d.U(0, 0.8)$, $\delta_{1i} \sim i.i.d.U(0, 0.5)$, and $\delta_{2i} \sim i.i.d.U(0, 0.2)$. Table 9 shows the good performance of the test when there exist two common breaks in the panels.

In summary, the size of our test is controlled for large N and T . The test exhibits monotonic power as the magnitude of the break increases and is powerful against various alternatives.

6. Empirical Example

In this section, we apply our approach to detect common breaks in the capital asset pricing model (CAPM). We use the Fama-French three-factor model augmented with the Carhart (1997) momentum factor considered in Antoch *et al.* (2019), which is given by

$$R_{it} - R_t^f = \alpha_{it} + \left(R_t^M - R_t^f\right) \beta_{it}^M + R_t^{HML} \beta_{it}^{HML} + R_t^{SMB} \beta_{it}^{SMB} + R_t^{MOM} \beta_{it}^{MOM} + u_{it},$$

for $1 \leq t \leq T$ and $1 \leq i \leq N$, where $R_{it} - R_t^f$ denotes the excess return on the mutual fund; the three factors include market risk premium, returns on a high minus low (HML) portfolio, and returns on a small minus big (SMB) portfolio; the momentum factor R_t^{MOM} describes the tendency of securities that have outperformed (or underperformed) the market over the past period to continue to outperform (or underperform) the market.

We test for common breaks in the coefficients for the mutual fund return data around the sub-prime crisis. Our sample period is from February 2005 to December 2011. Four factors can be downloaded from Ken French's data library.³ The monthly return data of mutual funds

³http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/data_library.html

are taken from Yahoo Finance. Mutual funds are classified according to their size, growth, value characteristics, and investment strategies. Using the Yahoo Finance classification, we focus on the characteristics of blend and growth to select ten categories of mutual funds. These are Foreign Large Blend, Foreign Small/Mid Blend, Foreign Large Growth, Foreign Small/Mid Growth, Large Blend, Mid-Cap Blend, Small Blend, Large Growth, Mid-Cap Growth, and Small Growth.

First, we apply the test to detect common breaks in the whole sample period of 2005M02–2011M12. The results in panel (a) of Table 10 show that the test rejects the one common break assumption at the 1% significance level for the nine categories. In this case, there are several possibilities such that there is no common break and each series (or some groups) has a distinct one or several breaks, or there are multiple common breaks. To see the overall tendency, we tentatively apply the Bai-Perron sequential test (Bai and Perron, 1998) to estimate the number and locations of the breaks for several mutual funds in each group. We find multiple break points, which are centralized at similar locations (early 2006, early 2008, and early 2009), even if the mutual funds are from different categories. These results suggest the possibility of multiple common breaks in mutual fund data.

Based on the above result and because Anotch *et al.* (2019) indicated that there exist structural changes in US mutual fund data during the sub-prime crisis period (mid-2008 to early 2009), we split the whole sample period into (b) the period before the sub-prime crisis (2005M02–2008M05), and (c) the period during the sub-prime crisis (2008M06–2011M12). In panel (b) of Table 10, the test rejects the null hypothesis for all categories in the period before the sub-prime crisis (before the middle of 2008); there still exists the possibility of distinct breaks in each series or multiple common breaks in this sub-period. On the other hand, as in panel (c) of Table 10, our test cannot reject the null hypothesis for the period 2008M06–2011M12. This result implies that the mutual fund data exhibit one common break during the sub-prime crisis.

7. Conclusion

In this study, we developed a new test based on the OLS residuals to detect whether structural breaks across individuals occurred at the common location in panel data models. The

asymptotic properties of the test were investigated under the null and alternative hypotheses. The simulation results indicated that the test is powerful against various alternatives. In application, we found evidence of the common break phenomenon in mutual fund data during the sub-prime crisis. Although we assumed cross-sectional independence throughout the paper, it may be interesting for the cross-sectional dependence in the error component to be generally taken into account. We leave such an extension for our future research.

References

- [1] Adesanya, O. (2020). Testing and dating structural breaks in generalized linear multivariate models for stock market contagion. Available at SSRN: <https://ssrn.com/abstract=3618001>.
- [2] Antoch, J., Hanousek, J., Horváth, L., Hušková, M., and Wang, S. X. (2019). Structural breaks in panel data: Large number of panels and short length time series. *Econometric Reviews*, 38(7), 828-855.
- [3] Bai, J. (1997). Estimation of a change point in multiple regression models. *Review of Economics and Statistics*, 79(4), 551-563.
- [4] Bai, J. (2010). Common breaks in means and variances for panel data. *Journal of Econometrics*, 157(1), 78-92.
- [5] Bai, J. and P. Perron (1998). Estimating and testing linear models with multiple structural changes. *Econometrica*, 66, 47-78.
- [6] Baltagi, B.H. Feng, Q. and Kao, C. (2016). Estimation of heterogeneous panels with structural breaks. *Journal of Econometrics*, 191(1), 176-195.
- [7] Baltagi, B. H., Kao, C., and Liu, L. (2017). Estimation and identification of change points in panel models with nonstationary or stationary regressors and error term. *Econometric Reviews*, 36(1-3), 85-102.
- [8] Billingsley P. (1968). Convergence of probability measures. Wiley. New York.

- [9] Carhart, M.M. (1997). On persistence in mutual fund performance. *The Journal of Finance*, 52, 57-82.
- [10] Chen, B., and Huang, L. (2018). Nonparametric testing for smooth structural changes in panel data models. *Journal of Econometrics*, 202(2), 245-267.
- [11] Claeys, P., and Vašíček, B. (2014). Measuring bilateral spillover and testing contagion on sovereign bond markets in Europe. *Journal of Banking & Finance*, 46, 151-165.
- [12] De Wachter, S., and Tzavalis, E. (2012). Detection of structural breaks in linear dynamic panel data models. *Computational Statistics & Data Analysis*, 56(11), 3020-3034.
- [13] Deng, A., and Perron, P. (2008). A non-local perspective on the power properties of the CUSUM and CUSUM of squares tests for structural change. *Journal of Econometrics*, 142, 212-240.
- [14] Hidalgo, J. and Schafgans, M. (2017). Inference and testing breaks in large dynamic panels with strong cross sectional dependence. *Journal of Econometrics*, 196(2), 259-274.
- [15] Horváth, L. and Hušková, M (2012). Change-point detection in panel data. *Journal of Time Series Analysis*, 33(4), 631-648.
- [16] Horváth, L. Hušková, M Rice, G., and Wang, J. (2017). Asymptotic properties of the cusum estimator for the time of change in linear panel data models. *Econometric Theory*, 33(2), 366-412.
- [17] Jiang, P., and Kurozumi, E. (2019). Power properties of the modified CUSUM tests. *Communications in Statistics - Theory and Methods*, 48(12), 2962-2981.
- [18] Kim, D. (2011). Estimating a common deterministic time trend break in large panels with cross sectional dependence. *Journal of Econometrics*, 164(2), 310-330.
- [19] Kim, D. (2014). Common breaks in time trends for large panel data with a factor structure. *Econometrics Journal*, 17(3), 301-337.

- [20] Li, D., Qian, J., and Su, L. (2016). Panel data models with interactive fixed effects and multiple structural breaks. *Journal of the American Statistical Association*, 111(516), 1804-1819.
- [21] Lumsdaine, R. L., Okui, R., and Wang, W. (2020). Estimation of panel group structure models with structural breaks in group memberships and coefficients. Available at SSRN: <https://ssrn.com/abstract=3617416>.
- [22] Oka, T. and Perron, P. (2018). Testing for common breaks in a multiple equations system. *Journal of Econometrics*, 204, 66-85.
- [23] Okui, R. and Wang, W. (2021). Heterogeneous structural breaks in panel data models. *Journal of Econometrics*, 220, 447-473.
- [24] Pauwels, L.L. Chan, F., and Griffoli, T.M. (2012). Testing for structural change in heterogeneous panels with an application to the Euro's trade effect. *Journal of Time Series Econometrics*, 4, 1-33.
- [25] Phillips, P.C.P. and Solo, V. (1992). Asymptotics for linear processes. *Annals of Statistics*, 20(2), 971-1001.
- [26] Qian, J. and Su, L. (2016). Shrinkage estimation of common breaks in panel data models via adaptive group fused lasso. *Journal of Econometrics*, 191, 86-109.
- [27] Shao, X., and Zhang, X., (2010). Testing for change points in time series. *Journal of the American Statistical Association*, 105(491), 1228-1240.
- [28] Vogelsang, T.J. (1999). Sources of nonmonotonic power when testing for a shift in mean of a dynamic time series. *Journal of Econometrics*, 88, 283-299.
- [29] Westerlund, J. (2019). Common breaks in means for cross-correlated fixed-T panel data. *Journal of Time Series Analysis*, 40(2), 248-255.
- [30] Yamazaki, D., and Kurozumi, E. (2015). Improving the finite sample performance of tests for a shift in mean. *Journal of Statistical Planning & Inference*, 40(2), 248-255.

Appendix A. Proof of Theorem 1

Supposing that the structural change occurred at a common location, Baltagi *et al.* (2016) showed the consistency of the common break estimator,

$$\lim_{(N,T) \rightarrow \infty} P(\hat{k} = k^0) = 1, \quad \text{which implies} \quad |\hat{k} - k^0| = o_p(1). \quad (\text{A.1})$$

In this Appendix, we derive the asymptotic distribution of the test statistic under the null hypothesis using this consistency property. We first focus on the limiting properties of the numerator of the statistic. Model (2) with the true common break k^0 is expressed as follows:

$$\begin{aligned} Y_i &= \bar{X}_i(k^0)b_i^0 + u_i \\ &= \bar{X}_i(\hat{k})b_i^0 + u_i + [\bar{X}_i(k^0) - \bar{X}_i(\hat{k})]b_i^0 \\ &= \bar{X}_i(\hat{k})b_i^0 + u_i + [Z_i(k^0) - Z_i(\hat{k})]\delta_i^0, \end{aligned} \quad (\text{A.2})$$

where $b_i^0 = [\beta_i^{0'}, \delta_i^{0'}]'$. Replacing Y_i with (A.2), the residuals in (4) can be rewritten as

$$\begin{aligned} \hat{u}_i &= \bar{X}_i(\hat{k})b_i^0 + u_i + [Z_i(k^0) - Z_i(\hat{k})]\delta_i^0 - \bar{X}_i(\hat{k})\hat{b}_i(\hat{k}) \\ &= u_i - \bar{X}_i(\hat{k})[\hat{b}_i(\hat{k}) - b_i^0] + [Z_i(k^0) - Z_i(\hat{k})]\delta_i^0 \\ &= u_i - X_i[\hat{\beta}_i(\hat{k}) - \beta_i^0] - Z_i(\hat{k})[\hat{\delta}_i(\hat{k}) - \delta_i^0] + [Z_i(k^0) - Z_i(\hat{k})]\delta_i^0, \end{aligned} \quad (\text{A.3})$$

whose vector form is represented by

$$\begin{bmatrix} \hat{u}_{i1} \\ \hat{u}_{i2} \\ \vdots \\ \hat{u}_{iT} \end{bmatrix} = \begin{bmatrix} u_{i1} \\ u_{i2} \\ \vdots \\ u_{iT} \end{bmatrix} - \begin{bmatrix} x_{i1} \\ x'_{i2} \\ \vdots \\ x'_{iT} \end{bmatrix} (\hat{\beta}_i(\hat{k}) - \beta_i^0) - \begin{bmatrix} 0 \\ \vdots \\ 0 \\ x'_{i(\hat{k}+1)} \\ \vdots \\ x'_{iT} \end{bmatrix} (\hat{\delta}_i(\hat{k}) - \delta_i^0) + \left(\begin{bmatrix} 0 \\ \vdots \\ 0 \\ x'_{i(k^0+1)} \\ \vdots \\ x'_{iT} \end{bmatrix} - \begin{bmatrix} 0 \\ \vdots \\ 0 \\ x'_{i(\hat{k}+1)} \\ \vdots \\ x'_{iT} \end{bmatrix} \right) \delta_i^0$$

For the sake of simplicity, \hat{k} is suppressed in $\hat{\beta}_i(\hat{k})$ and $\hat{\delta}_i(\hat{k})$. Then, the cumulative sum of

the residuals is

$$\begin{aligned}
& \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k \hat{u}_{it} \\
&= \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k x'_{it} (\hat{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^k x'_{it} (\hat{\delta}_i - \delta_i^0) 1_{\{k > \hat{k}\}} \\
&+ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=k^0+1}^k x'_{it} \delta_i^0 1_{\{k^0 < k \leq \hat{k}\}} + \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=k^0+1}^{\hat{k}} x'_{it} \delta_i^0 1_{\{k^0 < \hat{k} < k\}} \\
&- \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^k x'_{it} \delta_i^0 1_{\{\hat{k} < k \leq k^0\}} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^{k^0} x'_{it} \delta_i^0 1_{\{\hat{k} < k^0 < k\}} \\
&= U_1 - U_2 - U_3 + U_4 + U_5 - U_6 - U_7. \tag{A.4}
\end{aligned}$$

We can show that the terms U_4, U_5, U_6 , and U_7 are negligible as $N, T \rightarrow \infty$. Since k is bounded by k^0 and \hat{k} in U_4 , using the convergence property (A.1),

$$U_4 = \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=k^0+1}^k x'_{it} \delta_i^0 1_{\{k^0 < k \leq \hat{k}\}} = \sqrt{\frac{N}{T}} o_p(1) = o_p\left(\sqrt{\frac{N}{T}}\right). \tag{A.5}$$

Similarly, it is shown that the orders of terms U_5, U_6 , and U_7 are $o_p\left(\sqrt{\frac{N}{T}}\right)$, which will vanish since $N/T \rightarrow 0$ in Assumption 2(iii). The asymptotic distributions of the dominating terms U_1, U_2 , and U_3 are derived from Lemma A.1.

Lemma A.1 *Suppose that Assumptions 1–5 hold. We have, uniformly in $\tau \in (0, 1)$,*

$$\begin{aligned}
(i) & \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k u_{it} \Rightarrow \sigma W(\tau), \\
(ii) & \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k x'_{it} (\hat{\beta}_i - \beta_i^0) \Rightarrow \sigma \tau \frac{W(\tau^0)}{\tau^0}, \\
(iii) & \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^k x'_{it} (\hat{\delta}_i - \delta_i^0) 1_{\{k > \hat{k}\}} \Rightarrow \sigma(\tau - \tau^0) \left[\frac{W(1) - W(\tau^0)}{1 - \tau^0} - \frac{W(\tau^0)}{\tau^0} \right] 1_{\{\tau > \tau^0\}},
\end{aligned}$$

where $k = [T\tau]$, $k^0 = [T\tau^0]$, $W(\cdot)$ is a standard Brownian motion, and the long-run variance σ^2 is $\lim_{(N,T) \rightarrow \infty} E \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^T u_{it} \right)^2$.

Proof of Lemma A.1. (i) Denote the process

$$X_{N,T} \left(\frac{k}{T} \right) = \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k u_{it}.$$

It is shown that, for a particular τ ,

$$\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{[T\tau]} u_{it} \xrightarrow{d} \sigma W(\tau),$$

as $N, T \rightarrow \infty$. It remains to be shown that weak convergence holds uniformly in $\tau \in (0, 1)$. To this end, by Billingsley's (1968) Theorem 12.1, we next show that the moment condition (A.8) is satisfied such that the process $X_{N,T}(\tau)$ is tight. Applying Rosenthal's inequality, we have,

$$\begin{aligned} E \left| X_{N,T} \left(\frac{l}{T} \right) - X_{N,T} \left(\frac{k}{T} \right) \right|^{2\gamma} &= E \left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=k+1}^l u_{it} \right|^{2\gamma} \\ &\leq c_1 \sum_{i=1}^N E \left| \frac{1}{\sqrt{NT}} \sum_{t=k+1}^l u_{it} \right|^{2\gamma} + c_2 \left[\frac{1}{N} \sum_{i=1}^N E \left(\frac{1}{\sqrt{T}} \sum_{t=k+1}^l u_{it} \right)^2 \right]^\gamma \\ &\leq c_1 \sum_{i=1}^N E \left| \frac{1}{\sqrt{NT}} \sum_{t=k+1}^l u_{it} \right|^{2\gamma} + c_3 \left(\frac{l-k}{T} \right)^\gamma, \end{aligned} \tag{A.6}$$

with some constants c_1, c_2 , and c_3 . According to Phillips and Solo (1992) and p.637 of Horváth and Hušková (2012), the partial sum of u_{it} is composed of two parts,

$$\sum_{t=1}^k u_{it} = a_i \sum_{t=1}^k \epsilon_{it} + \eta_{ik},$$

where $\eta_{ik} = e_{i0}^* - e_{ik}^*$, $e_{it}^* = \sum_{l=1}^{\infty} c_{il}^* \epsilon_{i(t-l)}$, and $c_{il}^* = \sum_{k=l+1}^{\infty} c_{ik}$. For the term η_{ik} , Horváth and Hušková (2012, p.640) indicated that $E|\eta_{ik}|^\gamma \leq cE|\epsilon_{i0}|^\gamma$. Then, using Minkowski's inequality

and Rothenthal's inequality, we show that for $\gamma > 1$,

$$\begin{aligned}
E \left| \sum_{t=k+1}^l u_{it} \right|^{2\gamma} &= E \left| a_i \sum_{t=k+1}^l \epsilon_{it} + \eta_{il} - \eta_{ik} \right|^{2\gamma} \\
&\leq \left[\left(E \left| a_i \sum_{t=k+1}^l \epsilon_{it} \right|^{2\gamma} \right)^{\frac{1}{2\gamma}} + \left(E |\eta_{il} - \eta_{ik}|^{2\gamma} \right)^{\frac{1}{2\gamma}} \right]^{2\gamma} \\
&\leq \left\{ \left[c_4 \sum_{t=k+1}^l E |\epsilon_{it}|^{2\gamma} + c_5 \left(\sum_{t=k+1}^l E (\epsilon_{it})^2 \right)^\gamma \right]^{\frac{1}{2\gamma}} + \left(E |\eta_{il} - \eta_{ik}|^{2\gamma} \right)^{\frac{1}{2\gamma}} \right\}^{2\gamma} \\
&\leq \left\{ \left[c_4(l-k)E |\epsilon_{i0}|^{2\gamma} + c_5(l-k)^\gamma (E(\epsilon_{i0})^2)^\gamma \right]^{\frac{1}{2\gamma}} + \left(E |\eta_{il} - \eta_{ik}|^{2\gamma} \right)^{\frac{1}{2\gamma}} \right\}^{2\gamma} \\
&\leq \left\{ \left[c_6(l-k)^\gamma E |\epsilon_{i0}|^{2\gamma} \right]^{\frac{1}{2\gamma}} + \left(E |\epsilon_{i0}|^{2\gamma} \right)^{\frac{1}{2\gamma}} \right\}^{2\gamma} \\
&\leq c_7(l-k)^\gamma E |\epsilon_{i0}|^{2\gamma},
\end{aligned}$$

with some constants c_4 - c_7 . Then, we have, for $\gamma > 1$,

$$\begin{aligned}
\sum_{i=1}^N E \left| \frac{1}{\sqrt{NT}} \sum_{t=k+1}^l u_{it} \right|^{2\gamma} &= \frac{1}{(NT)^\gamma} \sum_{i=1}^N E \left| \sum_{t=k+1}^l u_{it} \right|^{2\gamma} \\
&\leq \frac{1}{(NT)^\gamma} \sum_{i=1}^N c_7(l-k)^\gamma E |\epsilon_{i0}|^{2\gamma} \\
&\leq c_7 \left(\frac{l-k}{T} \right)^\gamma \frac{1}{N} \sum_{i=1}^N E |\epsilon_{i0}|^{2\gamma} \\
&\leq c_8 \left(\frac{l-k}{T} \right)^\gamma, \tag{A.7}
\end{aligned}$$

with a constant c_8 . Combining (A.6) and (A.7), we can show that there exists constants $\gamma > 1$ and c_9 such that

$$E \left| X_{N,T} \left(\frac{l}{T} \right) - X_{N,T} \left(\frac{k}{T} \right) \right|^{2\gamma} \leq c_9 \left(\frac{l-k}{T} \right)^\gamma. \tag{A.8}$$

(ii) By regressing Y_i on $\bar{X}_i(\hat{k})$, the coefficient β_i is estimated as, if $\hat{k} \leq k^0$,

$$\hat{\beta}_i = \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} y_{it} = \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} (x'_{it} \beta_i^0 + u_{it}) = \beta_i^0 + \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it},$$

and if $\hat{k} > k^0$,

$$\begin{aligned}\hat{\beta}_i &= \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \left[\sum_{t=1}^{k^0} x_{it}(x'_{it}\beta_i^0 + u_{it}) + \sum_{t=k^0+1}^{\hat{k}} x_{it}(x'_{it}\beta_i^0 + x'_{it}\delta_i^0 + u_{it}) \right] \\ &= \beta_i^0 + \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}u_{it} + \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=k^0+1}^{\hat{k}} x_{it}x'_{it}\delta_i^0.\end{aligned}\quad (\text{A.9})$$

Then, we can see that,

$$\begin{aligned}\sqrt{T}(\hat{\beta}_i - \beta_i^0) &= \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{\hat{k}} x_{it}u_{it} + \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=k^0+1}^{\hat{k}} x_{it}x'_{it}\delta_i^0 1_{\{\hat{k} > k^0\}} \quad (\text{A.10}) \\ &= \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it}u_{it} + \left[\left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} - \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \right] \frac{1}{\sqrt{T}} \sum_{t=1}^{\hat{k}} x_{it}u_{it} \\ &\quad + \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \left[\frac{1}{\sqrt{T}} \sum_{t=1}^{\hat{k}} x_{it}u_{it} - \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it}u_{it} \right] + \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=k^0+1}^{\hat{k}} x_{it}x'_{it}\delta_i^0 1_{\{\hat{k} > k^0\}} \\ &= \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it}u_{it} + o_p\left(\frac{1}{T}\right) O_p(1) + O_p(1) o_p\left(\frac{1}{\sqrt{T}}\right) + o_p\left(\frac{1}{\sqrt{T}}\right) 1_{\{\hat{k} > k^0\}} \\ &= \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it}u_{it} + o_p\left(\frac{1}{T}\right) + o_p\left(\frac{1}{\sqrt{T}}\right),\end{aligned}\quad (\text{A.11})$$

where we replace \hat{k} with k^0 using the consistency property (A.1) and the following orders:

$$\begin{aligned}&\left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} - \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \\ &= \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} - \frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right) \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it}x'_{it} \right)^{-1} \\ &= O_p(1) o_p\left(\frac{1}{T}\right) O_p(1) = o_p\left(\frac{1}{T}\right), \\ &\quad \frac{1}{\sqrt{T}} \sum_{t=1}^{\hat{k}} x_{it}u_{it} - \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it}u_{it} = o_p\left(\frac{1}{\sqrt{T}}\right),\end{aligned}$$

and

$$\left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \frac{1}{T} \sum_{t=k^0+1}^{\hat{k}} x_{it}x'_{it}\delta_i^0 = O_p(1) o_p\left(\frac{1}{T}\right) = o_p\left(\frac{1}{T}\right).$$

Substituting (A.11) into the term U_2 in (A.4), we have

$$\begin{aligned}
& \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k x'_{it} (\hat{\beta}_i - \beta_i^0) = \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \sqrt{T} (\hat{\beta}_i - \beta_i^0) \\
& = \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} + \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(o_p \left(\frac{1}{T} \right) + o_p \left(\frac{1}{\sqrt{T}} \right) \right) \\
& = \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} + o_p \left(\frac{\sqrt{N}}{T} \right) + o_p \left(\sqrt{\frac{N}{T}} \right), \tag{A.12}
\end{aligned}$$

where the second and third terms in the last equality vanish since $N/T \rightarrow 0$ by Assumption 2(iii). From Assumptions 4(ii)–(iii), we can see that,

$$\left\| \frac{1}{k} \sum_{t=1}^k x'_{it} - c'_{i1} \right\| = o_p(1), \quad \text{and} \quad \left\| \left(\frac{1}{k} \sum_{t=1}^k x_{it} x'_{it} \right)^{-1} - C_i^{-1} \right\| = o_p(1). \tag{A.13}$$

Using orders in (A.13) and equality $c'_{i1} C_i^{-1} = [1, 0, \dots, 0]$, we have,

$$\begin{aligned}
& \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} - \frac{k}{k^0} \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{k^0} u_{it} \right| \\
& = \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} - \frac{k}{k^0} \frac{1}{\sqrt{N}} \sum_{i=1}^N c'_{i1} C_i^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} \right| \\
& \leq \frac{1}{\sqrt{N}} \sum_{i=1}^N \left\| \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} - \frac{k}{k^0} c'_{i1} C_i^{-1} \right\| \left\| \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} \right\| \\
& = \left\| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{k^0} x_{it} u_{it} \right\| o_p(1) = o_p(1). \tag{A.14}
\end{aligned}$$

Applying the functional central limit theorem (FCLT), we can see that

$$\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{k^0} u_{it} \Rightarrow \sigma W(\tau^0). \tag{A.15}$$

Hence, we have, uniformly in τ ,

$$\frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} \Rightarrow \sigma \tau \frac{W(\tau^0)}{\tau^0}, \tag{A.16}$$

(iii) The coefficient δ_i is estimated as, if $\hat{k} < k^0$,

$$\begin{aligned}
\hat{\delta}_i &= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}y_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}y_{it} \\
&= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \left[\sum_{t=\hat{k}+1}^{k^0} x_{it}(x'_{it}\beta_i^0 + u_{it}) + \sum_{t=k^0+1}^T x_{it}(x'_{it}\beta_i^0 + x'_{it}\delta_i^0 + u_{it}) \right] \\
&\quad - \left[\beta_i^0 + \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}u_{it} \right] \\
&= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}u_{it} + \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=k^0+1}^T x_{it}x'_{it}\delta_i^0 \\
&= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}u_{it} + \delta_i^0 - \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k^0} x_{it}x'_{it}\delta_i^0,
\end{aligned}$$

and if $\hat{k} \geq k^0$,

$$\begin{aligned}
\hat{\delta}_i &= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}y_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}y_{it} \\
&= \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}(x'_{it}\beta_i^0 + x'_{it}\delta_i^0 + u_{it}) \\
&\quad - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \left[\sum_{t=1}^{k^0} x_{it}(x'_{it}\beta_i^0 + u_{it}) + \sum_{t=k^0+1}^{\hat{k}} x_{it}(x'_{it}\beta_i^0 + x'_{it}\delta_i^0 + u_{it}) \right] \\
&= \delta_i^0 + \left(\sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it}u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it}u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \sum_{t=k^0+1}^{\hat{k}} x_{it}x'_{it}\delta_i^0.
\end{aligned}$$

Using the consistency property (A.1), the term $\sum_{t=\hat{k}+1}^{k^0} x_{it}x'_{it}\delta_i^0 = o_p(1)$ is negligible, and we can see that,

$$\sqrt{T}(\hat{\delta}_i - \delta_i^0) = \left(\frac{1}{T} \sum_{t=\hat{k}+1}^T x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=\hat{k}+1}^T x_{it}u_{it} - \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it}x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{\hat{k}} x_{it}u_{it} + o_p\left(\frac{1}{\sqrt{T}}\right). \tag{A.17}$$

Similar to the proof of (ii), \hat{k} in (A.17) can be replaced by k^0 due to the consistency of

$\hat{k} \xrightarrow{p} k^0$. Then, (A.17) is transformed into

$$\sqrt{T}(\hat{\delta}_i - \delta_i^0) = \left(\frac{1}{T} \sum_{t=k^0+1}^T x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=k^0+1}^T x_{it} u_{it} - \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} + o_p \left(\frac{1}{T} \right) + o_p \left(\frac{1}{\sqrt{T}} \right) \quad (\text{A.18})$$

Thus, we have,

$$\begin{aligned} & \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=k+1}^k x'_{it} (\hat{\delta}_i - \delta_i^0) = \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=k+1}^k x'_{it} \sqrt{T} (\hat{\delta}_i - \delta_i^0) \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k^0+1}^k x'_{it} 1_{\{k > k^0\}} + o_p \left(\frac{1}{T} \right) \right) \sqrt{T} (\hat{\delta}_i - \delta_i^0) \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k^0+1}^k x'_{it} 1_{\{k > k^0\}} \right) \sqrt{T} (\hat{\delta}_i - \delta_i^0) + o_p \left(\frac{\sqrt{N}}{T} \right) O_p(1) \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k^0+1}^k x'_{it} 1_{\{k > k^0\}} \right) \left[\left(\frac{1}{T} \sum_{t=k^0+1}^T x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=k^0+1}^T x_{it} u_{it} \right. \\ & \quad \left. - \left(\frac{1}{T} \sum_{t=1}^{k^0} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k^0} x_{it} u_{it} \right] + o_p \left(\frac{\sqrt{N}}{T} \right) + o_p \left(\sqrt{\frac{N}{T}} \right). \end{aligned} \quad (\text{A.19})$$

The terms in the brackets of (A.19) dominate the others. Similar to (A.14)–(A.16), we can see that, uniformly in $\tau \in (0, 1)$,

$$U_3 \Rightarrow \sigma(\tau - \tau^0) \left[\frac{W(1) - W(\tau^0)}{1 - \tau^0} - \frac{W(\tau^0)}{\tau^0} \right] 1_{\{\tau > \tau^0\}}.$$

Thus, we complete the proof of Lemma A.1. ■

Using (A.4), (A.5), and Lemma A.1, we can show that,

$$\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{[T\tau]} \hat{u}_{it} \Rightarrow \sigma W(\tau) - \sigma \tau \frac{W(\tau^0)}{\tau^0} - \sigma(\tau - \tau^0) \left[\frac{W(1) - W(\tau^0)}{1 - \tau^0} - \frac{W(\tau^0)}{\tau^0} \right] 1_{\{\tau > \tau^0\}},$$

uniformly in τ . Applying the continuous mapping theorem, we obtain,

$$\sup_{\tau \in \Omega(\epsilon)} \left| \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{[T\tau]} \hat{u}_{it} \right|^2 \Rightarrow \sup_{\tau \in \Omega(\epsilon)} \sigma^2 \left| W(\tau) - \tau \frac{W(\tau^0)}{\tau^0} - (\tau - \tau^0) \left[\frac{W(1) - W(\tau^0)}{1 - \tau^0} - \frac{W(\tau^0)}{\tau^0} \right] 1_{\{\tau > \tau^0\}} \right|^2. \quad (\text{A.20})$$

Next, we derive the asymptotic distribution of the normalization process under the null hypothesis. By definition (10), the normalization factor is based on the residuals \tilde{u}_{it} , which are calculated by regressing Y_i on $\tilde{X}_{it}(k_1, \hat{k}, k_2)$ in (8). We assume that $[T\epsilon] \leq k_1 \leq \hat{k} - [T\epsilon]$, and $\hat{k} + [T\epsilon] \leq k_2 \leq [T(1 - \epsilon)]$, where k_1, k_2 are bounded away from endpoints and the common break estimate \hat{k} . Since \hat{k} converges in probability to k^0 , we have $|k_j - \hat{k}| > |k^0 - \hat{k}|$ for $j = 1, 2$. Thus, we only consider the case in which k_1 and k_2 take values in $k_1 < k^0 < k_2$. In this case, the true model with a common break k^0 is written as

$$\begin{aligned}
Y_i &= [X_i, X_{1i}(k_1, k^0), X_{2i}(k^0, k_2), X_{3i}(k_2)] \begin{bmatrix} \beta_i^0 \\ 0 \\ \delta_i^0 \\ \delta_i^0 \end{bmatrix} + u_i \\
&= \tilde{X}_i(k_1, k^0, k_2) b_{1i}^0 + u_i \\
&= \tilde{X}_i(k_1, \hat{k}, k_2) b_{1i}^0 + u_i + [\tilde{X}_i(k_1, k^0, k_2) - \tilde{X}_i(k_1, \hat{k}, k_2)] b_{1i}^0. \tag{A.21}
\end{aligned}$$

The residuals are calculated by

$$\begin{aligned}
\tilde{u}_i &= \tilde{X}_i(k_1, \hat{k}, k_2) b_{1i}^0 + u_i + [\tilde{X}_i(k_1, k^0, k_2) - \tilde{X}_i(k_1, \hat{k}, k_2)] b_{1i}^0 - \tilde{X}_i(k_1, \hat{k}, k_2) \tilde{b}_{1i}(\hat{k}) \\
&= u_i + \tilde{X}_i(k_1, \hat{k}, k_2) b_{1i}^0 - \tilde{X}_i(k_1, \hat{k}, k_2) \tilde{b}_{1i}(\hat{k}) \\
&\quad + [0, X_{1i}(k_1, k^0) - X_{1i}(k_1, \hat{k}), X_{2i}(k^0, k_2) - X_{2i}(\hat{k}, k_2), 0] \begin{bmatrix} \beta_i^0 \\ 0 \\ \delta_i^0 \\ \delta_i^0 \end{bmatrix} \\
&= u_i - \tilde{X}_i(k_1, \hat{k}, k_2) (\tilde{b}_{1i}(\hat{k}) - b_{1i}^0) + [X_{2i}(k^0, k_2) - X_{2i}(\hat{k}, k_2)] \delta_i^0,
\end{aligned}$$

whose vector form is

$$\begin{aligned}
\begin{bmatrix} \tilde{u}_{i1} \\ \tilde{u}_{i2} \\ \vdots \\ \tilde{u}_{iT} \end{bmatrix} &= \begin{bmatrix} u_{i1} \\ u_{i2} \\ \vdots \\ u_{iT} \end{bmatrix} - \begin{bmatrix} x'_{i1} \\ x'_{i2} \\ \vdots \\ x'_{iT} \end{bmatrix} (\tilde{\beta}_i(\hat{k}) - \beta_i^0) - \begin{bmatrix} 0 \\ \vdots \\ 0 \\ x'_{i(k_1+1)} \\ \vdots \\ x'_{i\hat{k}} \\ 0 \\ \vdots \\ \vdots \\ 0 \end{bmatrix} \tilde{\delta}_{1i}(\hat{k}) - \begin{bmatrix} 0 \\ \vdots \\ \vdots \\ 0 \\ \vdots \\ x'_{i(\hat{k}+1)} \\ \vdots \\ x'_{ik_2} \\ 0 \\ \vdots \\ 0 \end{bmatrix} (\tilde{\delta}_{2i}(\hat{k}) - \delta_i^0) \\
&\quad - \begin{bmatrix} 0 \\ \vdots \\ \vdots \\ \vdots \\ 0 \\ x'_{i(k_2+1)} \\ \vdots \\ x'_{iT} \end{bmatrix} (\tilde{\delta}_{3i}(\hat{k}) - \delta_i^0) + \left(\begin{bmatrix} 0 \\ \vdots \\ \vdots \\ 0 \\ x'_{i(k^0+1)} \\ \vdots \\ x'_{ik_2} \\ 0 \\ \vdots \\ 0 \end{bmatrix} - \begin{bmatrix} 0 \\ \vdots \\ \vdots \\ 0 \\ \vdots \\ x'_{i(\hat{k}+1)} \\ \vdots \\ x'_{ik_2} \\ 0 \\ \vdots \\ 0 \end{bmatrix} \right) \delta_i^0. \tag{A.22}
\end{aligned}$$

For simplicity, \hat{k} is suppressed in $\tilde{\beta}_i(\hat{k})$, $\tilde{\delta}_{1i}(\hat{k})$, $\tilde{\delta}_{2i}(\hat{k})$, and $\tilde{\delta}_{3i}(\hat{k})$. The normalization factor is constructed by four terms V_1, V_2, V_3 , and V_4 , which are defined by

$$\begin{aligned}
V_1 &= \frac{1}{T} \sum_{s=1}^{k_1} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^s \tilde{u}_{it} \right)^2, \\
V_2 &= \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} \tilde{u}_{it} \right)^2, \\
V_3 &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s \tilde{u}_{it} \right)^2, \\
V_4 &= \frac{1}{T} \sum_{s=k_2+1}^T \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T \tilde{u}_{it} \right)^2.
\end{aligned}$$

Lemma A.2 derives the asymptotic distributions of the four terms under the null hypothesis.

Lemma A.2 *Suppose that Assumptions 1–5 hold. We have, as $N, T \rightarrow \infty$,*

$$\begin{aligned}
(i) \quad V_1 &\Rightarrow \sigma^2 \int_0^{\tau_1} \left(W(r) - \frac{r}{\tau_1} W(\tau_1) \right)^2 dr, \\
(ii) \quad V_2 &\Rightarrow \sigma^2 \int_{\tau_1}^{\tau^0} \left[W(\tau^0) - W(r) - \frac{\tau^0 - r}{\tau^0 - \tau_1} (W(\tau^0) - W(\tau_1)) \right]^2 dr, \\
(iii) \quad V_3 &\Rightarrow \sigma^2 \int_{\tau^0}^{\tau_2} \left[W(r) - W(\tau^0) - \frac{r - \tau^0}{\tau_2 - \tau^0} (W(\tau_2) - W(\tau^0)) \right]^2 dr, \\
(iv) \quad V_4 &\Rightarrow \sigma^2 \int_{\tau_2}^1 \left[W(1) - W(r) - \frac{1 - r}{1 - \tau_2} (W(1) - W(\tau_2)) \right]^2 dr.
\end{aligned}$$

Proof of Lemma A.2. (i) Using (A.22), the first term V_1 can be rewritten as

$$\begin{aligned}
V_1 &= \frac{1}{T} \sum_{s=1}^{k_1} \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \left[\sum_{t=1}^s u_{it} - \sum_{t=1}^s x'_{it} (\tilde{\beta}_i - \beta_i^0) \right] \right\}^2 \\
&= \frac{1}{T} \sum_{s=1}^{k_1} (V_{11} - V_{12})^2.
\end{aligned}$$

Using the FCLT, it is shown that

$$V_{11} \Rightarrow \sigma W(r). \quad (\text{A.23})$$

By the definition of $\tilde{\beta}_i$, we can see that,

$$\tilde{\beta}_i = \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} y_{it} = \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} (x'_{it} \beta_i^0 + u_{it}) = \beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it}.$$

Thus, we have,

$$\begin{aligned}
V_{12} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^s x'_{it} \left(\frac{1}{T} \sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k_1} x_{it} u_{it} \\
&\Rightarrow \sigma \frac{r}{\tau_1} W(\tau_1). \quad (\text{A.24})
\end{aligned}$$

Combining the results (A.23) and (A.24) and using the continuous mapping theorem, we can derive the asymptotic distribution of the first term V_1 as follows:

$$V_1 \Rightarrow \sigma^2 \int_0^{\tau_1} \left(W(r) - \frac{r}{\tau_1} W(\tau_1) \right)^2 dr.$$

(ii) The second term V_2 can be rewritten as

$$\begin{aligned}
V_2 &= \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \left[\sum_{t=s}^{\hat{k}} u_{it} - \sum_{t=s}^{\hat{k}} x'_{it}(\tilde{\beta}_i - \beta_i^0) - \sum_{t=s}^{\hat{k}} x'_{it}\tilde{\delta}_{1i} + \sum_{t=s}^{\hat{k}} x'_{it}\delta_i^0 1_{\{k^0 < s < \hat{k}\}} \right] \right\}^2 \\
&= \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} x'_{it}(\tilde{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} x'_{it}\tilde{\delta}_{1i} + o_p\left(\sqrt{\frac{N}{T}}\right) \right\}^2 \\
&= \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left(V_{21} - V_{22} - V_{23} + o_p\left(\sqrt{\frac{N}{T}}\right) \right)^2.
\end{aligned}$$

Since \hat{k} coincides asymptotically with the true break date from (A.1), \hat{k} in V_{21}, V_{22}, V_{23} can be replaced by k^0 . Then, we can show that

$$V_{21} = \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{\hat{k}} u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^{s-1} u_{it} \Rightarrow \sigma(W(\tau^0) - W(r)). \quad (\text{A.25})$$

$$\begin{aligned}
V_{22} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=s}^{\hat{k}} x'_{it} \sqrt{T}(\tilde{\beta}_i - \beta_i^0) \\
&= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=s}^{k^0} x'_{it} + o_p\left(\frac{1}{T}\right) \right) \left(\frac{1}{T} \sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k_1} x_{it} u_{it} \\
&\Rightarrow \sigma(\tau^0 - r) \frac{W(\tau_1)}{\tau_1}. \quad (\text{A.26})
\end{aligned}$$

The coefficient estimator $\tilde{\delta}_{1i}$ in V_{23} can be calculated as

$$\begin{aligned}
\tilde{\delta}_{1i} &= \left(\sum_{t=k_1+1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1+1}^{\hat{k}} x_{it} y_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} y_{it} \\
&= \left(\sum_{t=k_1+1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \left\{ \sum_{t=k_1+1}^{\hat{k}} x_{it} (x'_{it} \beta_i^0 + u_{it}) 1_{\{\hat{k} \leq k^0\}} + \left[\sum_{t=k_1+1}^{k^0} x_{it} (x'_{it} \beta_i^0 + u_{it}) \right. \right. \\
&\quad \left. \left. + \sum_{t=k^0+1}^{\hat{k}} x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) \right] 1_{\{\hat{k} > k^0\}} \right\} - \left[\beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\
&= \left(\sum_{t=k_1+1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1+1}^{\hat{k}} x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} + \left(\sum_{t=k_1+1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k^0\}} \\
&= \left(\sum_{t=k_1+1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1+1}^{\hat{k}} x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} + o_p\left(\frac{1}{T}\right).
\end{aligned}$$

Then, the third term V_{23} becomes, as $N, T \rightarrow \infty$,

$$\begin{aligned}
V_{23} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=s}^{\hat{k}} x'_{it} \sqrt{T} \tilde{\delta}_{1i} \\
&= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=s}^{k^0} x'_{it} + o_p \left(\frac{1}{T} \right) \right) \left[\sqrt{T} \left(\sum_{t=k_1+1}^{k^0} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1+1}^{k^0} x_{it} u_{it} - \sqrt{T} \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right. \\
&\quad \left. + o_p \left(\frac{1}{T} \right) + o_p \left(\frac{1}{\sqrt{T}} \right) + o_p \left(\frac{1}{\sqrt{T}} \right) \right] \\
&= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=s}^{k^0} x'_{it} \left[\sqrt{T} \left(\sum_{t=k_1+1}^{k^0} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1+1}^{k^0} x_{it} u_{it} - \sqrt{T} \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\
&\quad + o_p \left(\frac{\sqrt{N}}{\sqrt{T}} \right) + o_p \left(\frac{\sqrt{N}}{T} \right) + o_p \left(\frac{\sqrt{N}}{T^{3/2}} \right) + o_p \left(\frac{\sqrt{N}}{T^2} \right) \\
&\Rightarrow \sigma(\tau^0 - r) \left(\frac{W(\tau^0) - W(\tau_1)}{\tau^0 - \tau_1} - \frac{W(\tau_1)}{\tau_1} \right), \tag{A.27}
\end{aligned}$$

since $N/T \rightarrow 0$. Combining results (A.25), (A.26), and (A.27), we have

$$\begin{aligned}
V_2 &\Rightarrow \int_{\tau_1}^{\tau^0} \left[\sigma(W(\tau^0) - W(r)) - \sigma(\tau^0 - r) \frac{W(\tau_1)}{\tau_1} - \sigma(\tau^0 - r) \left(\frac{W(\tau^0) - W(\tau_1)}{\tau^0 - \tau_1} - \frac{W(\tau_1)}{\tau_1} \right) \right]^2 dr \\
&= \sigma^2 \int_{\tau_1}^{\tau^0} \left(W(\tau^0) - W(r) - (\tau^0 - r) \frac{W(\tau^0) - W(\tau_1)}{\tau^0 - \tau_1} \right)^2 dr.
\end{aligned}$$

(iii) The third term V_3 can be rewritten as

$$\begin{aligned}
V_3 &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \left[\sum_{t=\hat{k}+1}^s u_{it} - \sum_{t=\hat{k}+1}^s x'_{it} (\tilde{\beta}_i - \beta_i^0) - \sum_{t=\hat{k}+1}^s x'_{it} (\tilde{\delta}_{2i} - \delta_i^0) \right. \right. \\
&\quad \left. \left. - \sum_{t=\hat{k}+1}^{k^0} x'_{it} \delta_i^0 1_{\{\hat{k} < k^0 < s\}} - \sum_{t=\hat{k}+1}^s x'_{it} \delta_i^0 1_{\{s \leq k^0\}} \right] \right\}^2 \\
&= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left[\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s x'_{it} (\tilde{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s x'_{it} (\tilde{\delta}_{2i} - \delta_i^0) \right. \\
&\quad \left. - o_p \left(\sqrt{\frac{N}{T}} \right) \right]^2 \\
&= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2} \left(V_{31} - V_{32} - V_{33} - o_p \left(\sqrt{\frac{N}{T}} \right) \right)^2.
\end{aligned}$$

Similar to (A.25) and (A.27), we can find that

$$V_{31} \Rightarrow \sigma(W(r) - W(\tau^0)), \quad (\text{A.28})$$

$$\begin{aligned} V_{32} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k^0+1}^s x'_{it} + o_p\left(\frac{1}{T}\right) \right) \left(\frac{1}{T} \sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k_1} x_{it} u_{it} \\ &\Rightarrow \sigma(r - \tau^0) \frac{W(\tau_1)}{\tau_1}. \end{aligned} \quad (\text{A.29})$$

The coefficient δ_{2i} is estimated as

$$\begin{aligned} \tilde{\delta}_{2i} &= \left(\sum_{t=\hat{k}+1}^{k_2} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2} x_{it} y_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} y_{it} \\ &= \left(\sum_{t=\hat{k}+1}^{k_2} x_{it} x'_{it} \right)^{-1} \left\{ \sum_{t=\hat{k}+1}^{k_2} x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) 1_{\{k^0 \leq \hat{k}\}} + \left[\sum_{t=\hat{k}+1}^{k^0} x_{it} (x'_{it} \beta_i^0 + u_{it}) \right. \right. \\ &\quad \left. \left. + \sum_{t=k^0+1}^{k_2} x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) \right] 1_{\{k^0 < \hat{k}\}} \right\} - \left[\beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &= \delta_i^0 + \left(\sum_{t=\hat{k}+1}^{k_2} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2} x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} - o_p\left(\frac{1}{T}\right). \end{aligned}$$

Then, similar to V_{23} , the term V_{33} becomes, as $N, T \rightarrow \infty$,

$$\begin{aligned} &\frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=\hat{k}+1}^s x'_{it} \sqrt{T} (\tilde{\delta}_{2i} - \delta_i^0) \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k^0+1}^s x'_{it} + o_p\left(\frac{1}{T}\right) \right) \left[\sqrt{T} \left(\sum_{t=k^0+1}^{k_2} x_{it} x'_{it} \right)^{-1} \sum_{t=k^0+1}^{k_2} x_{it} u_{it} - \sqrt{T} \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right. \\ &\quad \left. + o_p\left(\frac{1}{T}\right) + o_p\left(\frac{1}{\sqrt{T}}\right) + o_p\left(\frac{1}{\sqrt{T}}\right) \right] \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=k^0+1}^s x'_{it} \left[\sqrt{T} \left(\sum_{t=k^0+1}^{k_2} x_{it} x'_{it} \right)^{-1} \sum_{t=k^0+1}^{k_2} x_{it} u_{it} - \sqrt{T} \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &\quad + o_p\left(\frac{\sqrt{N}}{\sqrt{T}}\right) + o_p\left(\frac{\sqrt{N}}{T}\right) + o_p\left(\frac{\sqrt{N}}{T^{3/2}}\right) + o_p\left(\frac{\sqrt{N}}{T^2}\right) \\ &\Rightarrow \sigma(r - \tau^0) \left(\frac{W(\tau_2) - W(\tau^0)}{\tau_2 - \tau^0} - \frac{W(\tau_1)}{\tau_1} \right), \end{aligned} \quad (\text{A.30})$$

since $N/T \rightarrow 0$. Combining results (A.28), (A.29), and (A.30), we have

$$V_3 \Rightarrow \sigma^2 \int_{\tau^0}^{\tau_2} \left(W(r) - W(\tau^0) - (r - \tau^0) \frac{W(\tau_2) - W(\tau^0)}{\tau_2 - \tau^0} \right)^2 dr.$$

(iv) The fourth term V_4 can be rewritten as

$$\begin{aligned} V_4 &= \frac{1}{T} \sum_{s=k_2+1}^T \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \left[\sum_{t=s}^T u_{it} - \sum_{t=s}^T x'_{it}(\tilde{\beta}_i - \beta_i^0) - \sum_{t=s}^T x'_{it}(\tilde{\delta}_{3i} - \delta_i^0) \right] \right\}^2 \\ &= \frac{1}{T} \sum_{s=k_2+1}^T \left[\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T u_{it} x'_{it}(\tilde{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T u_{it} x'_{it}(\tilde{\delta}_{3i} - \delta_i^0) \right]^2 \\ &= \frac{1}{T} \sum_{s=k_2+1}^T (V_{41} - V_{42} - V_{43})^2. \end{aligned}$$

It is easily seen that

$$V_{41} \Rightarrow \sigma(W(1) - W(r)), \quad (\text{A.31})$$

$$\begin{aligned} V_{42} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=s}^T x'_{it} \right) \left(\frac{1}{T} \sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=1}^{k_1} x_{it} u_{it} \\ &\Rightarrow \sigma(1-r) \frac{W(\tau_1)}{\tau_1}. \end{aligned} \quad (\text{A.32})$$

The coefficient estimator $\tilde{\delta}_{3i}$ can be written as

$$\begin{aligned} \tilde{\delta}_{3i} &= \left(\sum_{t=k_2+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=k_2+1}^T x_{it} y_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} y_{it} \\ &= \left(\sum_{t=k_2+1}^T x_{it} x'_{it} \right)^{-1} \left[\sum_{t=k_2+1}^T x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) \right] - \left[\beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &= \delta_i^0 + \left(\sum_{t=k_2+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=k_2+1}^T x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it}. \end{aligned}$$

Then, it is shown that, as $N, T \rightarrow \infty$,

$$\begin{aligned} V_{43} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=s}^T x'_{it} \sqrt{T} (\tilde{\delta}_{3i} - \delta_i^0) \\ &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=s}^T x'_{it} \right) \sqrt{T} \left[\left(\sum_{t=k_2+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=k_2+1}^T x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &\Rightarrow \sigma(1-r) \left(\frac{W(1) - W(\tau_2)}{1 - \tau_2} - \frac{W(\tau_1)}{\tau_1} \right), \end{aligned} \quad (\text{A.33})$$

since $N/T \rightarrow 0$. From (A.31), (A.32), and (A.33), we have

$$V_4 \Rightarrow \sigma^2 \int_{\tau_2}^1 \left[W(1) - W(r) - \frac{1-r}{1-\tau_2} (W(1) - W(\tau_2)) \right]^2 dr.$$

The proof of Lemma A.2 is complete. ■

Proof of Theorem 1. Combining the asymptotic distributions in (A.20) and Lemma A.2, we can complete the proof. ■

Appendix B. Proof of Theorem 2

We first investigate the statistical properties of the common break estimator \hat{k} under the alternative hypothesis in Proposition 1. The proof follows the proof of Lemmas 1–2 and Theorem 1 in Baltagi *et al.* (2016). Suppose that there are two groups and the individuals in the same group share a common break date. These groups are denoted by $G_1 = \{i : \text{individuals in group 1 with a common break } k_1^0\}$ and $G_2 = \{i : \text{individuals in group 2 with common break } k_2^0\}$. The model under the alternative can be specified as

$$\begin{cases} y_{it} = x'_{it}\beta_i^0 + x'_{it}\delta_i^0 1_{(t>k_1^0)} + u_{it} & t = 1, \dots, T, \text{ for } i \in G_1, \\ y_{it} = x'_{it}\beta_i^0 + x'_{it}\delta_i^0 1_{(t>k_2^0)} + u_{it} & t = 1, \dots, T, \text{ for } i \in G_2. \end{cases}$$

The vector form can be rewritten as

$$Y_i = [X_i, Z_i(k_j^0)]b_i^0 + u_i, \text{ for } i \in G_j, j = 1, 2.$$

The common break point is estimated in (3) by minimizing the total sum of the squared OLS residuals. Let SSR_i denote the sum of the squared residuals of regression Y_i on X_i (no break case). Using the equality on page 185 of Baltagi *et al.* (2016),

$$SSR_i - SSR_i(k^*) = \hat{\delta}_i(k^*)' [Z_i(k^*)' M_i Z_i(k^*)] \hat{\delta}_i(k^*),$$

estimation (3) can be transformed into

$$\begin{aligned}
\hat{k} &= \arg \min_{1 \leq k^* \leq T-1} \sum_{i=1}^N SSR_i(k^*) \\
&= \arg \max_{1 \leq k^* \leq T-1} \sum_{i=1}^N (SSR_i - SSR_i(k^*)) \\
&= \arg \max_{1 \leq k^* \leq T-1} \sum_{i=1}^N SV_i(k^*) \\
&= \arg \max_{1 \leq k^* \leq T-1} \left[\sum_{i \in G_1} (SV_i(k^*) - SV_i(k_1^0)) + \sum_{i \in G_2} (SV_i(k^*) - SV_i(k_1^0)) \right], \quad (\text{B.1})
\end{aligned}$$

where

$$\begin{aligned}
M_i &= I - X_i(X_i'X_i)^{-1}X_i', \\
SV_i(k^*) &= \hat{\delta}_i(k^*)'[Z_i(k^*)'M_iZ_i(k^*)]\hat{\delta}_i(k^*), \\
SV_i(k_j^0) &= \hat{\delta}_i(k_j^0)'[Z_i(k_j^0)'M_iZ_i(k_j^0)]\hat{\delta}_i(k_j^0), \text{ for } j = 1, 2.
\end{aligned}$$

For individuals in group 1, we can see that the coefficient estimators are given by

$$\begin{aligned}
\hat{\delta}_i(k^*) &= [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)M_iY_i, \\
\hat{\delta}_i(k_1^0) &= [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)M_iY_i.
\end{aligned}$$

Replacing Y_i with

$$Y_i = X_i\beta_i^0 + Z_i(k_1^0)\delta_i^0 + u_i,$$

we have

$$\begin{aligned}
\hat{\delta}_i(k^*) &= [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_i[X_i\beta_i^0 + Z_i(k_1^0)\delta_i^0 + u_i] \\
&= [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_iZ_i(k_1^0)\delta_i^0 + [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_iu_i, \\
\hat{\delta}_i(k_1^0) &= [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)'M_i[X_i\beta_i^0 + Z_i(k_1^0)\delta_i^0 + u_i] \\
&= \delta_i^0 + [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)'M_iu_i.
\end{aligned}$$

Similarly, for individuals in group 2, by replacing Y_i with

$$Y_i = X_i\beta_i^0 + Z_i(k_2^0)\delta_i^0 + u_i,$$

the coefficients estimators are rewritten as

$$\begin{aligned}
\hat{\delta}_i(k^*) &= [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_i[X_i\beta_i^0 + Z_i(k_2^0)\delta_i^0 + u_i] \\
&= [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_iZ_i(k_2^0)\delta_i^0 + [Z_i(k^*)'M_iZ_i(k^*)]^{-1}Z_i(k^*)'M_iu_i, \\
\hat{\delta}_i(k_1^0) &= [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)'M_i[X_i\beta_i^0 + Z_i(k_2^0)\delta_i^0 + u_i] \\
&= [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)'M_iZ_i(k_2^0)\delta_i^0 + [Z_i(k_1^0)'M_iZ_i(k_1^0)]^{-1}Z_i(k_1^0)'M_iu_i.
\end{aligned}$$

To simplify the notations, we use Z_i, Z_{1i}^0, Z_{2i}^0 to replace $Z_i(k^*), Z_i(k_1^0), Z_i(k_2^0)$. For the individuals in group 1, we have

$$\begin{aligned}
SV_i(k^*) &= \delta_i^{0'}Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iZ_{1i}^0\delta_i^0 + 2\delta_i^{0'}Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i \\
&\quad + u_i'M_iZ_i'(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i, \tag{B.2}
\end{aligned}$$

$$SV_i(k_1^0) = \delta_i^{0'}Z_{1i}^{0'}M_iZ_{1i}^0\delta_i^0 + 2\delta_i^{0'}Z_{1i}^{0'}M_iu_i + u_i'M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^{0'}M_iu_i. \tag{B.3}$$

Using (B.2) and (B.3), $SV_i(k^*) - SV_i(k_1^0)$ becomes

$$\begin{aligned}
SV_i(k^*) - SV_i(k_1^0) &= -\delta_i^{0'} \left[Z_{1i}^{0'}M_iZ_{1i}^0 - Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iZ_{1i}^0 \right] \delta_i^0 \\
&\quad + 2\delta_i^{0'}Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i - 2\delta_i^{0'}Z_{1i}^{0'}M_iu_i \\
&\quad + u_i'M_iZ_i'(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i - u_i'M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^{0'}M_iu_i,
\end{aligned}$$

and can be decomposed into the term defined by

$$J_{1i}(k^*) = \delta_i^{0'} \left[Z_{1i}^{0'}M_iZ_{1i}^0 - Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iZ_{1i}^0 \right] \delta_i^0 \tag{B.4}$$

and the term related to disturbance u_i defined by

$$\begin{aligned}
H_{1i}(k^*) &= 2\delta_i^{0'}Z_{1i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i - 2\delta_i^{0'}Z_{1i}^{0'}M_iu_i \\
&\quad + u_i'M_iZ_i'(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i - u_i'M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^{0'}M_iu_i. \tag{B.5}
\end{aligned}$$

Then, we have $SV_i(k^*) - SV_i(k_1^0) = -J_{1i}(k^*) + H_{1i}(k^*)$ for $i \in G_1$. A similar transformation for individuals in group 2 shows that

$$\begin{aligned}
SV_i(k^*) &= \delta_i^{0'}Z_{2i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iZ_{2i}^0\delta_i^0 + 2\delta_i^{0'}Z_{2i}^{0'}M_iZ_i(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i \\
&\quad + u_i'M_iZ_i'(Z_i'M_iZ_i)^{-1}Z_i'M_iu_i, \tag{B.6}
\end{aligned}$$

$$\begin{aligned}
SV_i(k_1^0) &= \delta_i^{0'}Z_{2i}^{0'}M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^0M_iZ_{2i}^0\delta_i^0 + 2\delta_i^{0'}Z_{2i}^{0'}M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^0M_iu_i \\
&\quad + u_i'M_iZ_{1i}^0(Z_{1i}^{0'}M_iZ_{1i}^0)^{-1}Z_{1i}^0M_iu_i. \tag{B.7}
\end{aligned}$$

Using (B.6) and (B.7), we can see that, for $i \in G_2$,

$$SV_i(k^*) - SV_i(k_1^0) = -J_{2i}(k^*) + H_{2i}(k^*),$$

where the term $J_{2i}(k^*)$ is denoted by

$$J_{2i}(k^*) = \delta_i^{0'} \left[Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i Z_{2i}^0 - Z_{2i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i Z_{2i}^0 \right] \delta_i^0 \quad (\text{B.8})$$

and the term $H_{2i}(k^*)$ related to disturbance is denoted by

$$\begin{aligned} H_{2i}(k^*) &= 2\delta_i^{0'} Z_{2i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - 2\delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i u_i \\ &\quad + u_i' M_i Z_i' (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - u_i' M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^{0'} M_i u_i. \end{aligned} \quad (\text{B.9})$$

Thus, (B.1) can be rewritten as

$$\hat{k} = \arg \max_{1 \leq k^* \leq T-1} \left[\sum_{i \in G_1} (-J_{1i}(k^*) + H_{1i}(k^*)) + \sum_{i \in G_2} (-J_{2i}(k^*) + H_{2i}(k^*)) \right].$$

Define the sets $K(C_1) = \{k : 1 \leq k < k_1^0 - C_1\}$, $K(C_2) = \{k : k_2^0 + C_2 < k \leq T\}$, and $K(C) = K(C_1) \cup K(C_2) = \{k : 1 \leq k < k_1^0 - C_1 \text{ or } k_2^0 + C_2 < k \leq T\}$ with positive constants C_1, C_2 . Next, we show that the common break estimator cannot appear in the set $K(C_1)$ by Lemmas B.1–B.2. A similar result can be obtained for the set $K(C_2)$ by symmetry; thus, the details are omitted. Define

$$Z_{1i}^\Delta = \begin{cases} Z_i(k^*) - Z_i(k_1^0) & \text{if } k^* < k_1^0 \\ -(Z_i(k^*) - Z_i(k_1^0)) & \text{if } k^* \geq k_1^0 \end{cases} \text{ and } Z_{2i}^\Delta = \begin{cases} Z_i(k^*) - Z_i(k_2^0) & \text{if } k^* < k_2^0 \\ -(Z_i(k^*) - Z_i(k_2^0)) & \text{if } k^* \geq k_2^0 \end{cases}.$$

Lemma B.1 *Under Assumptions 1–6, for all large N and T , with probability tending to 1,*

$$\inf_{k^* \in K(C_1)} \frac{1}{k_1^0 - k^*} \left(\sum_{i \in G_1} J_{1i}(k^*) + \sum_{i \in G_2} J_{2i}(k^*) \right) \geq \lambda \phi_N.$$

Proof of Lemma B.1. We first show that the summation of part $J_{1i}(k^*)$ has a lower bound in the case of $k^* \in K(C_1)$. From Lemma A.2 in Bai(1997), if $k^* < k_1^0$,

$$\begin{aligned} J_{1i}(k^*) &= \delta_i^{0'} \left[Z_{1i}^{0'} M_i Z_{1i}^0 - Z_{1i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i Z_{1i}^0 \right] \delta_i^0 \\ &= \delta_i^{0'} Z_{1i}^{\Delta'} Z_{1i}^\Delta (Z_i' Z_i)^{-1} Z_{1i}^{0'} Z_{1i}^0 \delta_i^0. \end{aligned} \quad (\text{B.10})$$

Since the matrix

$$\frac{Z_{1i}^{\Delta'} Z_{1i}^{\Delta}}{k_1^0 - k^*} \left(\frac{Z_i' Z_i}{T} \right)^{-1} \frac{Z_{1i}^{0'} Z_{1i}^0}{T} \quad (\text{B.11})$$

is symmetric and positive definite from Assumption 4, we have

$$\frac{1}{k_1^0 - k^*} \sum_{i \in G_1} J_{1i}(k^*) = \sum_{i \in G_1} \delta_i^{0'} S_i' \Lambda_i S_i \delta_i^0 = \sum_{i \in G_1} \tilde{\delta}_i^{0'} \Lambda_i \tilde{\delta}_i^0 \geq \sum_{i \in G_1} \lambda_i \tilde{\delta}_i^{0'} \tilde{\delta}_i^0, \quad (\text{B.12})$$

where Λ_i is a diagonal matrix comprising of the eigenvalues of matrix (B.11), $\tilde{\delta}_i^0 = S_i \delta_i^0$, and λ_i is the minimum eigenvalue of (B.11). Since $\tilde{\delta}_i^{0'} \tilde{\delta}_i^0 = \delta_i^{0'} S_i' S_i \delta_i^0 = \delta_i^{0'} \delta_i^0$, with probability tending to one for large N and T , we show that

$$\frac{1}{k_1^0 - k^*} \sum_{i \in G_1} J_{1i}(k^*) \geq \lambda_1 \sum_{i \in G_1} \delta_i^{0'} \delta_i^0 = \lambda_1 \phi_{N_1}, \quad (\text{B.13})$$

where $\lambda_1 = \min_{i \in G_1} \{\lambda_i\}$. Next, we investigate the lower bound of $J_{2i}(k^*)$ for individuals in group 2. Denote

$$V_i(a, b) = \begin{bmatrix} x'_{i(a+1)} \\ x'_{i(a+2)} \\ \vdots \\ x'_{ib} \end{bmatrix}, \quad V_i^0(a, b, c) = \begin{bmatrix} 0^{(b-a) \times p} \\ x'_{i(b+1)} \\ x'_{i(b+2)} \\ \vdots \\ x'_{ic} \end{bmatrix}, \quad \text{and} \quad S = \begin{bmatrix} I & 0 \\ -I & I \end{bmatrix},$$

where $V_i(a, b)$ is a $(b - a) \times p$ matrix whose j th row is the same as the $(a + j)$ th row of X_i , $V_i^0(a, b, c)$ is a $(c - a) \times p$ matrix whose first $b - a$ rows are zeros and the j th row is the same as the $(a + j)$ th row of X_i for $j > b - a$, and S is a $2p \times 2p$ matrix constructed by $p \times p$ identity matrix I . The second term $J_{2i}(k)$ can be transformed into

$$\begin{aligned} J_{2i}(k^*) &= \delta_i^{0'} \left[Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i Z_{2i}^0 - Z_{2i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i Z_{2i}^0 \right] \delta_i^0 \\ &= \delta_i^{0'} Z_{2i}^{0'} \left[M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i - M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i \right] Z_{2i}^0 \delta_i^0 \\ &= \delta_i^{0'} Z_{2i}^{0'} \left[M_i - M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i - \left(M_i - M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i \right) \right] Z_{2i}^0 \delta_i^0 \\ &= \delta_i^{0'} Z_{2i}^{0'} \left(M_W - M_{W_1^0} \right) Z_{2i}^0 \delta_i^0 \\ &= \delta_i^{0'} Z_{2i}^{0'} \left(M_{\bar{W}} - M_{\bar{W}_1^0} \right) Z_{2i}^0 \delta_i^0, \end{aligned} \quad (\text{B.14})$$

where

$$\begin{aligned}
W &= [X_i, Z_i(k^*)] = \begin{bmatrix} V_i(0, k^*) & 0 \\ V_i(k^*, T) & V_i(k^*, T) \end{bmatrix}, \quad W_1^0 = [X_i, Z_i(k_1^0)] = \begin{bmatrix} V_i(0, k_1^0) & 0 \\ V_i(k_1^0, T_1^0) & V_i(k_1^0, T) \end{bmatrix}, \\
\bar{W} &= \begin{bmatrix} V_i(0, k^*) & 0 \\ 0 & V_i(k^*, T) \end{bmatrix}, \quad \bar{W}_1^0 = \begin{bmatrix} V_i(0, k_1^0) & 0 \\ 0 & V_i(k_1^0, T) \end{bmatrix}, \\
M_X &= I - X(X'X)^{-1}X', \text{ for matrix } X.
\end{aligned}$$

The final equality (B.14) holds because

$$\begin{aligned}
M_W &= I - W(W'W)^{-1}W' = I - WS(S'W'WS)^{-1}S'W' = I - \bar{W}(\bar{W}'\bar{W})^{-1}\bar{W} = M_{\bar{W}}, \\
M_{W_1^0} &= I - W_1^0(W_1^{0'}W_1^0)^{-1}W_1^{0'} = I - W_1^0S(S'W_1^{0'}W_1^0S)^{-1}S'W_1^{0'} = I - \bar{W}_1^0(\bar{W}_1^{0'}\bar{W}_1^0)^{-1}\bar{W}_1^{0'} = M_{\bar{W}_1^0}.
\end{aligned}$$

Since \bar{W} and \bar{W}_1^0 are block matrices, it follows that

$$\begin{aligned}
& Z_{2i}^{0'} M_{\bar{W}} Z_{2i}^0 \\
&= Z_{2i}^{0'} \left[I - \bar{W} (\bar{W}'\bar{W})^{-1} \bar{W} \right] Z_{2i}^0 \\
&= [0, V_i^0(k^*, k_2^0, T)]' \left[I - \bar{W} (\bar{W}'\bar{W})^{-1} \bar{W} \right] \begin{bmatrix} 0 \\ V_i^0(k^*, k_2^0, T) \end{bmatrix} \\
&= V_i^0(k^*, k_2^0, T)' V_i^0(k^*, k_2^0, T) - V_i^0(k^*, k_2^0, T)' V_i(k^*, T) (V_i(k^*, T)' V_i(k^*, T))^{-1} V_i(k^*, T) V_i^0(k^*, k_2^0, T) \\
&= V_i(k_2^0, T)' V_i(k_2^0, T) - V_i(k_2^0, T)' V_i(k_2^0, T) (V_i(k^*, T)' V_i(k^*, T))^{-1} V_i(k_2^0, T)' V_i(k_2^0, T) \\
&= V_i(k_2^0, T)' V_i(k_2^0, T) \left[\left(V_i(k_2^0, T)' V_i(k_2^0, T) \right)^{-1} - \left(V_i(k^*, T)' V_i(k^*, T) \right)^{-1} \right] V_i(k_2^0, T)' V_i(k_2^0, T), \quad (\text{B.15})
\end{aligned}$$

$$\begin{aligned}
& Z_{2i}^{0'} M_{\bar{W}_1^0} Z_{2i}^0 \\
&= Z_{2i}^{0'} \left[I - \bar{W}_1^0 (\bar{W}_1^{0'}\bar{W}_1^0)^{-1} \bar{W}_1^{0'} \right] Z_{2i}^0 \\
&= [0, V_i^0(k_1^0, k_2^0, T)]' \left[I - \bar{W}_1^0 (\bar{W}_1^{0'}\bar{W}_1^0)^{-1} \bar{W}_1^{0'} \right] \begin{bmatrix} 0 \\ V_i^0(k_1^0, k_2^0, T) \end{bmatrix} \\
&= V_i^0(k_1^0, k_2^0, T)' V_i^0(k_1^0, k_2^0, T) - V_i^0(k_1^0, k_2^0, T)' V_i(k_1^0, T) (V_i(k_1^0, T)' V_i(k_1^0, T))^{-1} V_i(k_1^0, T) V_i^0(k_1^0, k_2^0, T) \\
&= V_i(k_2^0, T)' V_i(k_2^0, T) \left[\left(V_i(k_2^0, T)' V_i(k_2^0, T) \right)^{-1} - \left(V_i(k_1^0, T)' V_i(k_1^0, T) \right)^{-1} \right] V_i(k_2^0, T)' V_i(k_2^0, T). \quad (\text{B.16})
\end{aligned}$$

Substituting (B.15) and (B.16) into (B.14), we have

$$\begin{aligned}
J_{2i}(k^*) &= \delta_i^{0'} V_i(k_2^0, T)' V_i(k_2^0, T) \left[\left(V_i(k_1^0, T)' V_i(k_1^0, T) \right)^{-1} - \left(V_i(k^*, T)' V_i(k^*, T) \right)^{-1} \right] V_i(k_2^0, T)' V_i(k_2^0, T) \delta_i^0 \\
&= \delta_i^{0'} Z_{2i}^{0'} Z_{2i}^0 \left[\left(Z_{1i}^{0'} Z_{1i}^0 \right)^{-1} - \left(Z_i' Z_i \right)^{-1} \right] Z_{2i}^{0'} Z_{2i}^0 \delta_i^0 \\
&= \delta_i^{0'} Z_{2i}^{0'} Z_{2i}^0 \left(Z_i' Z_i \right)^{-1} \left(Z_i' Z_i - Z_{1i}^{0'} Z_{1i}^0 \right) \left(Z_{1i}^{0'} Z_{1i}^0 \right)^{-1} Z_{2i}^{0'} Z_{2i}^0 \delta_i^0 \\
&= \delta_i^{0'} Z_{2i}^{0'} Z_{2i}^0 \left(Z_i' Z_i \right)^{-1} Z_{1i}^{\Delta'} Z_{1i}^{\Delta} \left(Z_{1i}^{0'} Z_{1i}^0 \right)^{-1} Z_{2i}^{0'} Z_{2i}^0 \delta_i^0,
\end{aligned} \tag{B.17}$$

which is symmetric from the first equality. Hence, under Assumptions 4 and 5,

$$\frac{Z_{2i}^{0'} Z_{2i}^0}{T} \left(\frac{Z_i' Z_i}{T} \right)^{-1} \frac{Z_{1i}^{\Delta'} Z_{1i}^{\Delta}}{k_1^0 - k^*} \left(\frac{Z_{1i}^{0'} Z_{1i}^0}{T} \right)^{-1} \frac{Z_{2i}^{0'} Z_{2i}^0}{T} \tag{B.18}$$

is positive definite. Then, we have

$$\frac{1}{k_1^0 - k^*} \sum_{i \in G_2} J_{2i}(k^*) \geq \lambda_2 \sum_{i \in G_2} \delta_i^{0'} \delta_i^0 = \lambda_2 \phi_{N_2}, \tag{B.19}$$

where $\lambda_2 = \min_{i \in G_2} \{\lambda_i\}$, and λ_i is the minimum eigenvalue of matrix (B.18). From inequalities (B.13) and (B.19), the proof of Lemma B.1 is complete. ■

Lemma B.2 Under Assumptions 1–6, uniformly on $k^* \in K(C_1)$,

$$\begin{aligned}
(i) \quad & \sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} u_i = O_p(\sqrt{\phi_{N_1}}), \\
(ii) \quad & \sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i = O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right), \\
(iii) \quad & \sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i = O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right), \\
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^{\Delta} (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i = O_p\left(\sqrt{\frac{\phi_{N_2}}{T}}\right), \\
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i = O_p\left(\sqrt{\phi_{N_2}}\right) + O_p\left(\sqrt{\frac{\phi_{N_2}}{T}}\right), \\
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 \left[(Z_i' M_i Z_i)^{-1} - (Z_{1i}^0' M_i Z_{1i}^0)^{-1} \right] Z_{1i}^{\Delta'} M_i u_i = O_p\left(\sqrt{\frac{\phi_{N_2}}{T}}\right), \\
(iv) \quad & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^{\Delta} (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i = O_p\left(\frac{N}{T}\right), \\
(v) \quad & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^{\Delta} (Z_i' M_i Z_i)^{-1} Z_{1i}^0' M_i u_i = O_p\left(\frac{N}{T}\right) + O_p\left(\frac{N}{\sqrt{T}}\right), \\
(vi) \quad & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^0 \left[(Z_i' M_i Z_i)^{-1} - (Z_{1i}^0' M_i Z_{1i}^0)^{-1} \right] Z_{1i}^0' M_i u_i = O_p\left(\frac{N}{T}\right).
\end{aligned}$$

Proof of Lemma B.2. (i) It is shown that

$$\frac{1}{\sqrt{k_1^0 - k^*}} Z_{1i}^{\Delta'} u_i = O_p(1), \quad \text{since} \quad \text{Var}\left(\frac{1}{\sqrt{k_1^0 - k^*}} Z_{1i}^{\Delta'} u_i\right) < \infty.$$

Then, we have

$$\sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} u_i = O_p(\sqrt{\phi_{N_1}}).$$

(ii) We can show that

$$\sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i = \frac{1}{\sqrt{T}} \sum_{i \in G_1} \delta_i^{0'} \frac{Z_{1i}^{\Delta'} X_i}{k_1^0 - k^*} \left(\frac{X_i' X_i}{T}\right)^{-1} \frac{1}{\sqrt{T}} X_i' u_i = O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right),$$

since for large T

$$\frac{1}{\sqrt{T}} X_i' u_i = O_p(1).$$

(iii) By expanding M_i , we can show that

$$\begin{aligned}
& \sum_{i \in G_1} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{1i}^{\Delta'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \\
&= \frac{1}{\sqrt{T}} \sum_{i \in G_1} \delta_i^{0'} \frac{Z_{1i}^{\Delta'} Z_i}{k_1^0 - k^*} \left(\frac{Z_i' M_i Z_i}{T} \right)^{-1} \frac{1}{\sqrt{T}} Z_i' M_i u_i \\
&\quad - \frac{1}{\sqrt{T}} \sum_{i \in G_1} \delta_i^{0'} \frac{Z_{1i}^{\Delta'} X_i}{k_1^0 - k^*} \left(\frac{X_i' X_i}{T} \right)^{-1} \frac{X_i' Z_i}{T} \left(\frac{Z_i' M_i Z_i}{T} \right)^{-1} \frac{1}{\sqrt{T}} Z_i' M_i u_i \\
&= O_p \left(\sqrt{\frac{\phi_{N_1}}{T}} \right).
\end{aligned}$$

To prove the second order, since $Z_{2i}^{0'} Z_{1i}^{\Delta} = 0$, we have

$$\begin{aligned}
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^{\Delta} (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \\
&= \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} Z_{1i}^{\Delta} (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \\
&\quad - \frac{1}{\sqrt{T}} \sum_{i \in G_2} \delta_i^{0'} \frac{Z_{2i}^{0'} X_i}{T} \left(\frac{X_i' X_i}{T} \right)^{-1} \frac{X_i' Z_{1i}^{\Delta}}{k_1^0 - k^*} \left(\frac{Z_i' M_i Z_i}{T} \right)^{-1} \frac{1}{\sqrt{T}} Z_i' M_i u_i \\
&= O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right).
\end{aligned}$$

Considering the third order, we can show that

$$\begin{aligned}
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i \\
&= \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} u_i \\
&\quad - \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i \\
&= \frac{1}{\sqrt{k_1^0 - k^*}} O_p \left(\sqrt{\phi_{N_2}} \right) + O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right) \\
&= O_p \left(\sqrt{\phi_{N_2}} \right) + O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right).
\end{aligned}$$

The last term can be transformed into

$$\begin{aligned}
& \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} \left(Z_{1i}^0 M_i Z_{1i}^0 - Z_i' M_i Z_i \right) \left(Z_{1i}^0 M_i Z_{1i}^0 \right)^{-1} Z_{1i}^{0'} M_i u_i \\
&= \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} \left(-Z_{1i}^{\Delta'} Z_{1i}^{\Delta} - Z_{1i}^{\Delta'} M_i Z_{1i}^0 - Z_{1i}^0 M_i Z_{1i}^{\Delta} \right) \left(Z_{1i}^0 M_i Z_{1i}^0 \right)^{-1} Z_{1i}^{0'} M_i u_i \\
&\quad + \sum_{i \in G_2} \frac{1}{k_1^0 - k^*} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} \left(Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' Z_{1i}^{\Delta} \right) \left(Z_{1i}^0 M_i Z_{1i}^0 \right)^{-1} Z_{1i}^{0'} M_i u_i \\
&= O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right) + O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right).
\end{aligned}$$

(iv) The term $\sum_{i=1}^N \frac{1}{T(k_1^0 - k^*)} u_i' M_i Z_{1i}^{\Delta} \left(\frac{Z_i' M_i Z_i}{T} \right)^{-1} Z_{1i}^{\Delta'} M_i u_i$ has the same order as that of $\sum_{i=1}^N \frac{1}{T(k_1^0 - k^*)} u_i' M_i Z_{1i}^{\Delta} Z_{1i}^{\Delta'} M_i u_i$ since the matrix $\frac{Z_i' M_i Z_i}{T} = O_p(1)$ for large T. Expanding matrix M_i , we have

$$\begin{aligned}
& \sum_{i=1}^N \frac{1}{T(k_1^0 - k^*)} u_i' M_i Z_{1i}^{\Delta} Z_{1i}^{\Delta'} M_i u_i \\
&= \frac{1}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} u_i - \frac{2}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' X_i (X_i' X_i)^{-1} X_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} u_i \\
&\quad + \frac{1}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' X_i (X_i' X_i)^{-1} X_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i \tag{B.20}
\end{aligned}$$

Consider the first term in (B.20),

$$\frac{1}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} u_i = O_p \left(\frac{N}{T} \right).$$

Similarly, it can be shown that the second term,

$$\begin{aligned}
& \frac{1}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' X_i (X_i' X_i)^{-1} X_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} u_i \\
&= \sqrt{\frac{k_1^0 - k^*}{T}} \frac{1}{T} \sum_{i=1}^N \frac{u_i' X_i}{\sqrt{T}} \left(\frac{X_i' X_i}{T} \right)^{-1} \frac{X_i' Z_{1i}^{\Delta}}{k_1^0 - k^*} \frac{Z_{1i}^{\Delta'} u_i}{\sqrt{k_1^0 - k^*}} = O_p \left(\frac{N}{T} \right),
\end{aligned}$$

and the third term,

$$\begin{aligned}
& \frac{1}{T(k_1^0 - k^*)} \sum_{i=1}^N u_i' X_i (X_i' X_i)^{-1} X_i' Z_{1i}^{\Delta} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i \\
&= \frac{k_1^0 - k^*}{T} \frac{1}{T} \sum_{i=1}^N \frac{u_i' X_i}{\sqrt{T}} \left(\frac{X_i' X_i}{T} \right)^{-1} \frac{X_i' Z_{1i}^{\Delta}}{k_1^0 - k^*} \frac{Z_{1i}^{\Delta'} X_i}{k_1^0 - k^*} \left(\frac{X_i' X_i}{T} \right)^{-1} \frac{X_i' u_i}{\sqrt{T}} = O_p \left(\frac{N}{T} \right).
\end{aligned}$$

Thus, we have

$$\sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i = O_p \left(\frac{N}{T} \right).$$

(v) By expanding M_i , we show that

$$\begin{aligned} & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_{1i}^{0'} M_i u_i \\ = & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_{1i}^{0'} M_i u_i - \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' X_i (X_i' X_i)^{-1} X_i' Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_{1i}^{0'} M_i u_i \\ = & O_p \left(\frac{N}{\sqrt{T}} \right) + O_p \left(\frac{N}{T} \right). \end{aligned}$$

(vi) We show that

$$\begin{aligned} & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} \left(Z_{1i}^{0'} M_i Z_{1i}^0 - Z_i' M_i Z_i \right) \left(Z_{1i}^{0'} M_i Z_{1i}^0 \right)^{-1} Z_{1i}^{0'} M_i u_i \\ = & \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} \left(-Z_{1i}^{\Delta'} M_i Z_{1i}^\Delta - Z_{1i}^{\Delta'} M_i Z_{1i}^0 - Z_{1i}^{0'} M_i Z_{1i}^\Delta \right) \left(Z_{1i}^{0'} M_i Z_{1i}^0 \right)^{-1} Z_{1i}^{0'} M_i u_i \\ = & O_p \left(\frac{N}{T} \right). \end{aligned}$$

The proof of Lemma B.2 is complete. ■

Proof of Proposition 1. We first show that for any given $\epsilon > 0$,

$$P \left(\sup_{K(C_1)} \left| \frac{\sum_{i \in G_1} H_{1i}(k^*) + \sum_{i \in G_2} H_{2i}(k^*)}{k_1^0 - k^*} \right| \geq \lambda \phi_N \right) < \epsilon. \quad (\text{B.21})$$

Using (B.5) and (B.9), we see that the sum of $H_{1i}(k^*) + H_{2i}(k^*)$ can be decomposed into

three parts:

$$\begin{aligned}
& \frac{1}{k_1^0 - k^*} \left(\sum_{i \in G_1} H_{1i}(k^*) + \sum_{i \in G_2} H_{2i}(k^*) \right) \\
&= \frac{1}{k_1^0 - k^*} \left[\sum_{i \in G_1} 2\delta_i^{0'} Z_{1i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - \sum_{i \in G_1} 2\delta_i^{0'} Z_{1i}^{0'} M_i u_i \right] \\
&+ \frac{1}{k_1^0 - k^*} \left[\sum_{i \in G_2} 2\delta_i^{0'} Z_{2i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - \sum_{i \in G_2} 2\delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i u_i \right] \\
&+ \frac{1}{k_1^0 - k^*} \left[\sum_{i=1}^N u_i' M_i Z_i' (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - \sum_{i=1}^N u_i' M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^0 M_i u_i \right] \\
&= H_1 + H_2 + H_3.
\end{aligned}$$

Consider the first term by replacing Z_{1i}^0 with $Z_i - Z_{1i}^\Delta$,

$$\begin{aligned}
|H_1| &= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \left[\delta_i^{0'} Z_{1i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - \delta_i^{0'} Z_{1i}^{0'} M_i u_i \right] \right| \\
&= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \left[\delta_i^{0'} Z_i' M_i u_i - \delta_i^{0'} Z_{1i}^{\Delta'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - \delta_i^{0'} Z_i' M_i u_i + \delta_i^{0'} Z_{1i}^{\Delta'} M_i u_i \right] \right| \\
&= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \left[\delta_i^{0'} Z_{1i}^{\Delta'} M_i u_i - \delta_i^{0'} Z_{1i}^{\Delta'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \right] \right| \\
&\leq 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \delta_i^{0'} Z_{1i}^{\Delta'} u_i \right| + 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \delta_i^{0'} Z_{1i}^{\Delta'} X_i (X_i' X_i)^{-1} X_i' u_i \right| \\
&\quad + 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_1} \delta_i^{0'} Z_{1i}^{\Delta'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \right| \\
&= O_p(\sqrt{\phi_{N_1}}) + O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right) + O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right), \tag{B.22}
\end{aligned}$$

where the inequality is obtained by expanding M_i , and the final equality uses the orders in (i)–(iii) of Lemma B.2.

For the second term H_2 , replacing Z_i with $Z_{1i}^\Delta + Z_{1i}^0$, and using (iii) of Lemma B.2,

$$\begin{aligned}
|H_2| &= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \left[2\delta_i^{0'} Z_{2i}^{0'} M_i Z_i (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - 2\delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^{0'} M_i u_i \right] \right| \\
&= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \left[\delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i + \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \right. \right. \\
&\quad \left. \left. - \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^{0'} M_i u_i \right] \right| \\
&= 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \left[\delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i + \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i \right. \right. \\
&\quad \left. \left. + \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 \left[(Z_i' M_i Z_i)^{-1} - (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} \right] Z_{1i}^{0'} M_i u_i \right] \right| \\
&\leq 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^\Delta (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i \right| \\
&\quad + 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i \right| \tag{B.23}
\end{aligned}$$

$$\begin{aligned}
&\quad + 2 \left| \frac{1}{k_1^0 - k^*} \sum_{i \in G_2} \delta_i^{0'} Z_{2i}^{0'} M_i Z_{1i}^0 \left[(Z_i' M_i Z_i)^{-1} - (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} \right] Z_{1i}^{0'} M_i u_i \right| \\
&= O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right) + O_p(\sqrt{\phi_{N_2}}) + O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right) + O_p \left(\sqrt{\frac{\phi_{N_2}}{T}} \right). \tag{B.24}
\end{aligned}$$

By (iv)–(vi) of Lemma B.2, the order of the third term is

$$\begin{aligned}
|H_3| &= \left| \frac{1}{k_1^0 - k^*} \sum_{i=1}^N \left[u_i' M_i Z_i' (Z_i' M_i Z_i)^{-1} Z_i' M_i u_i - u_i' M_i Z_{1i}^0 (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} Z_{1i}^{0'} M_i u_i \right] \right| \\
&\leq \left| \frac{1}{k_1^0 - k^*} \sum_{i=1}^N u_i' M_i Z_{1i}^{\Delta'} (Z_i' M_i Z_i)^{-1} Z_{1i}^{\Delta'} M_i u_i \right| + \left| \frac{2}{k_1^0 - k^*} \sum_{i=1}^N u_i' M_i Z_{1i}^{\Delta'} (Z_i' M_i Z_i)^{-1} Z_{1i}^{0'} M_i u_i \right| \\
&\quad + \left| \sum_{i=1}^N \frac{1}{k_1^0 - k^*} u_i' M_i Z_{1i}^0 \left[(Z_i' M_i Z_i)^{-1} - (Z_{1i}^{0'} M_i Z_{1i}^0)^{-1} \right] Z_{1i}^{0'} M_i u_i \right| \\
&= O_p \left(\frac{N}{T} \right) + O_p \left(\frac{N}{T} \right) + O_p \left(\frac{N}{\sqrt{T}} \right) + O_p \left(\frac{N}{T} \right). \tag{B.25}
\end{aligned}$$

Combining (B.22), (B.24), and (B.25) under Assumption 2, the term,

$$\begin{aligned} & \frac{1}{\phi_N} \left| \frac{\sum_{i \in G_1} H_{1i}(k^*) + \sum_{i \in G_2} H_{2i}(k^*)}{k_1^0 - k^*} \right| \\ &= \frac{1}{\phi_N} \left| \left[O_p(\sqrt{\phi_{N_1}}) + O_p(\sqrt{\phi_{N_2}}) + O_p\left(\sqrt{\frac{\phi_{N_1}}{T}}\right) + O_p\left(\sqrt{\frac{\phi_{N_2}}{T}}\right) + O_p\left(\frac{N}{\sqrt{T}}\right) + O_p\left(\frac{N}{T}\right) \right] \right| \\ &\rightarrow 0, \end{aligned}$$

will vanish for any $k^* \in K(C_1)$. On the other hand, the part $\phi_N^{-1}(k_1^0 - k^*)^{-1} \left| \sum_{i \in G_1} J_{1i}(k^*) + \sum_{i \in G_2} J_{2i}(k^*) \right|$ has a lower bound from Lemma B.1. Hence, for any $\epsilon > 0$,

$$P \left(\sup_{K(C_1)} \left| \frac{\sum_{i \in G_1} H_{1i}(k^*) + \sum_{i \in G_2} H_{2i}(k^*)}{k_1^0 - k^*} \right| \geq \sup_{K(C_1)} \left| \frac{\sum_{i \in G_1} J_{1i}(k^*) + \sum_{i \in G_2} J_{2i}(k^*)}{k_1^0 - k^*} \right| \right) < \epsilon,$$

which implies that

$$\begin{aligned} P \left(\sup_{K(C_1)} \sum_{i \in G_1} -J_{1i}(k^*) + H_{1i}(k^*) + \sum_{i \in G_2} -J_{2i}(k^*) + H_{2i}(k^*) \geq 0 \right) &< \epsilon, \\ P \left(\sup_{K(C_1)} \sum_{i=1}^N [SV_i(k^*) - SV_i(k_1^0)] \geq 0 \right) &< \epsilon. \end{aligned}$$

Finally, we obtain that for any given $\epsilon > 0$, and both large N and T ,

$$P(\hat{k} \in K(C_1)) < \epsilon.$$

In other words, the total sum of squared residuals cannot be maximized in the case of $k^* \in K(C_1)$. By symmetry, the estimation of the common break point (3) can be transformed into

$$\hat{k} = \arg \max_{1 \leq k^* \leq T-1} \left[\sum_{i \in G_1} (SV_i(k^*) - SV_i(k_2^0)) + \sum_{i \in G_2} (SV_i(k^*) - SV_i(k_2^0)) \right].$$

Similarly, we can show that, for any given $\epsilon > 0$,

$$P(\hat{k} \in K(C_2)) < \epsilon.$$

The common break point estimator is obtained in set $K(C_2)$ with probability tending to zero.

Thus, we complete the proof of Proposition 1. ■

Proposition 1 indicates that the estimated common break is stochastically bounded by either true break points or located between k_1^0 and k_2^0 . Then, we can say that

$$\frac{k_1^0 - \hat{k}}{T} = O_p\left(\frac{1}{T}\right), \quad \text{if } \hat{k} \leq k_1^0, \quad (\text{B.26})$$

$$\frac{\hat{k} - k_2^0}{T} = O_p\left(\frac{1}{T}\right), \quad \text{if } \hat{k} \geq k_2^0. \quad (\text{B.27})$$

Using this property of the common break estimator under the alternative, we next show that the numerator of the statistic will diverge under H_{1A} .

Proof of Proposition 2. Under the alternative, from (A.4), the CUSUM of the residuals for individuals in group j ($j = 1, 2$) are calculated as

$$\begin{aligned} & \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=1}^k \hat{u}_{it} \\ &= \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=1}^k u_{it} - \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=1}^k x'_{it} (\hat{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=\hat{k}+1}^k x'_{it} (\hat{\delta}_i - \delta_i^0) 1_{\{k > \hat{k}\}} \\ &+ \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=k_j^0+1}^k x'_{it} \delta_i^0 1_{\{k_j^0 < k \leq \hat{k}\}} + \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=k_j^0+1}^{\hat{k}} x'_{it} \delta_i^0 1_{\{k_j^0 < \hat{k} < k\}} \\ &- \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=\hat{k}+1}^k x'_{it} \delta_i^0 1_{\{\hat{k} < k \leq k_j^0\}} - \frac{1}{\sqrt{NT}} \sum_{i \in G_j} \sum_{t=\hat{k}+1}^{k_j^0} x'_{it} \delta_i^0 1_{\{\hat{k} < k_j^0 < k\}}. \end{aligned}$$

Then, the total sum of the squared residuals $\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k \hat{u}_{it}$ is expressed as

$$\begin{aligned}
& \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k x'_{it} (\hat{\beta}_i(\hat{k}) - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^k x'_{it} (\hat{\delta}_i(\hat{k}) - \delta_i^0) 1_{\{k > \hat{k}\}} \\
& + \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=k_1^0+1}^k x'_{it} \delta_i^0 1_{\{k_1^0 < k \leq \hat{k}\}} + \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=k_1^0+1}^{\hat{k}} x'_{it} \delta_i^0 1_{\{k_1^0 < \hat{k} < k\}} \\
& - \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=\hat{k}+1}^k x'_{it} \delta_i^0 1_{\{\hat{k} < k \leq k_1^0\}} - \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=\hat{k}+1}^{k_1^0} x'_{it} \delta_i^0 1_{\{\hat{k} < k_1^0 < k\}} \\
& + \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=k_2^0+1}^k x'_{it} \delta_i^0 1_{\{k_2^0 < k \leq \hat{k}\}} + \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=k_2^0+1}^{\hat{k}} x'_{it} \delta_i^0 1_{\{k_2^0 < \hat{k} < k\}} \\
& - \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^k x'_{it} \delta_i^0 1_{\{\hat{k} < k \leq k_2^0\}} - \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^{k_2^0} x'_{it} \delta_i^0 1_{\{\hat{k} < k_2^0 < k\}} \\
& = U_1^{H_1} - U_2^{H_1} - U_3^{H_1} + U_4^{H_1} + U_5^{H_1} - U_6^{H_1} - U_7^{H_1} + U_8^{H_1} + U_9^{H_1} - U_{10}^{H_1} - U_{11}^{H_1}. \quad (\text{B.28})
\end{aligned}$$

Since $\hat{k} < k_1^0$ in $U_6^{H_1}, U_7^{H_1}$, and $\hat{k} > k_2^0$ in $U_8^{H_1}, U_9^{H_1}$, using the orders (B.26) and (B.27), we have

$$U_6^{H_1} = O_p \left(\sqrt{\frac{N}{T}} \right), U_7^{H_1} = O_p \left(\sqrt{\frac{N}{T}} \right), U_8^{H_1} = O_p \left(\sqrt{\frac{N}{T}} \right), U_9^{H_1} = O_p \left(\sqrt{\frac{N}{T}} \right). \quad (\text{B.29})$$

From (A.10), we know that for $i \in G_j, j = 1, 2$,

$$\begin{aligned}
\sqrt{T}(\hat{\beta}_i - \beta_i^0) &= \sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + \sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_j^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_j^0\}} \\
&= \begin{cases} \sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + \sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_1^0\}} & \text{if } j = 1, \\ \sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + O_p \left(\frac{1}{\sqrt{T}} \right) & \text{if } j = 2, \end{cases}
\end{aligned}$$

using order (B.27). Then, the second term $U_2^{H_1}$ becomes

$$\begin{aligned}
& \frac{1}{\sqrt{N}} \sum_{i \in G_1} \frac{1}{T} \sum_{t=1}^k x'_{it} \sqrt{T} \left[\left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_1^0\}} \right] \\
& + \frac{1}{\sqrt{N}} \sum_{i \in G_2} \frac{1}{T} \sum_{t=1}^k x'_{it} \left[\sqrt{T} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + O_p \left(\frac{1}{\sqrt{T}} \right) \right] \\
& = O_p(1) + \frac{1}{\sqrt{N}} \sum_{i \in G_1} \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_1^0\}} \\
& + O_p(1) + O_p \left(\sqrt{\frac{N}{T}} \right) \\
& = O_p(1) + U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} + O_p(1) + O_p \left(\sqrt{\frac{N}{T}} \right). \tag{B.30}
\end{aligned}$$

Considering the third term $U_3^{H_1}$, for individuals $i \in G_j$, the coefficient estimator is

$$\begin{aligned}
\hat{\delta}_i & = \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it} y_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} y_{it} \\
& = \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \left\{ \sum_{t=\hat{k}+1}^T x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) 1_{\{\hat{k} \geq k_j^0\}} \right. \\
& + \left. \left[\sum_{t=\hat{k}+1}^{k_j^0} x_{it} (x'_{it} \beta_i^0 + u_{it}) + \sum_{t=k_j^0+1}^T x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) \right] 1_{\{\hat{k} < k_j^0\}} \right\} \\
& - \left[\beta_i^0 + \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} + \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_j^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_j^0\}} \right] \\
& = \delta_i^0 + \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it} u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} \\
& - \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_j^0} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} < k_j^0\}} - \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_j^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_j^0\}},
\end{aligned}$$

where the fourth term in the final equality is $O_p(1/T)$ for individuals in group 1 using order (B.26), while the fifth term is $O_p(1/T)$ for individuals in group 2 using order (B.27). Then,

the third term $U_3^{H_1}$ can be rewritten as

$$\begin{aligned}
& \left\{ \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=\hat{k}+1}^k x'_{it} \sqrt{T} \left[\left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^T x_{it} u_{it} - \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{\hat{k}} x_{it} u_{it} \right] \right. \\
& - O_p \left(\sqrt{\frac{N}{T}} \right) - \frac{1}{\sqrt{N}} \sum_{i \in G_1} \frac{1}{T} \sum_{t=\hat{k}+1}^k x'_{it} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} > k_1^0\}} \\
& \left. - \frac{1}{\sqrt{N}} \sum_{i \in G_2} \frac{1}{T} \sum_{t=\hat{k}+1}^k x'_{it} \left(\frac{1}{T} \sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \frac{1}{\sqrt{T}} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \delta_i^0 1_{\{\hat{k} < k_2^0\}} - O_p \left(\sqrt{\frac{N}{T}} \right) \right\} 1_{\{k > \hat{k}\}} \\
& = \left[O_p(1) - O_p \left(\sqrt{\frac{N}{T}} \right) - U_{31}^{H_1} 1_{\{\hat{k} > k_1^0\}} - U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} - O_p \left(\sqrt{\frac{N}{T}} \right) \right] 1_{\{k > \hat{k}\}}. \tag{B.31}
\end{aligned}$$

Thus, from (B.29), (B.30), and (B.31), (B.28) can be rewritten as

$$\begin{aligned}
& \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k \hat{u}_{it} \\
& = O_p(1) - \left[O_p(1) + U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} + O_p(1) + O_p \left(\sqrt{\frac{N}{T}} \right) \right] \\
& - \left[O_p(1) - O_p \left(\sqrt{\frac{N}{T}} \right) - U_{31}^{H_1} 1_{\{\hat{k} > k_1^0\}} - U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} - O_p \left(\sqrt{\frac{N}{T}} \right) \right] 1_{\{k > \hat{k}\}} \\
& + U_4^{H_1} + U_5^{H_1} + O_p \left(\sqrt{\frac{N}{T}} \right) - U_{10}^{H_1} - U_{11}^{H_1} \\
& = -U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} + \left[U_{31}^{H_1} 1_{\{\hat{k} > k_1^0\}} + U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} \right] 1_{\{k > \hat{k}\}} + U_4^{H_1} + U_5^{H_1} - U_{10}^{H_1} - U_{11}^{H_1} \\
& + O_p(1) + O_p \left(\sqrt{\frac{N}{T}} \right). \tag{B.32}
\end{aligned}$$

Next, we show that (B.32) diverges at the rate of \sqrt{NT} under the alternative in the following three cases.

Case (i). Suppose that $\hat{k} < k_1^0 < k_2^0$, we have

$$U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} = 0, U_{31}^{H_1} 1_{\{\hat{k} > k_1^0\}} = 0, U_4^{H_1} = 0, U_5^{H_1} = 0.$$

Choosing $k \in (k_1^0 + C_1, k_2^0]$, we can see that $U_{11}^{H_1} = 0$, and

$$\begin{aligned}
& -U_{10}^{H_1} + U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} 1_{\{k > \hat{k}\}} \\
&= -\frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^k x'_{it} \delta_i^0 + \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^k x'_{it} \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \delta_i^0 \\
&= -\sqrt{\frac{T}{N}} \sum_{i \in G_2} \frac{1}{T} \sum_{t=\hat{k}+1}^k x'_{it} \left(\frac{1}{T} \sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \frac{1}{T} \sum_{t=k_2^0+1}^T x_{it} x'_{it} \delta_i^0 \\
&= O_p(\sqrt{NT}).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\sup_{k \in [1, T-1]} US_{NT}(k, \hat{k}) &\geq \sup_{k \in (k_1^0 + C_1, k_2^0]} US_{NT}(k, \hat{k}) \\
&= \sup_{k \in (k_1^0 + C_1, k_2^0]} \left(U_{32}^{H_1} - U_{10}^{H_1} + O_p(1) + O_p\left(\sqrt{\frac{N}{T}}\right) \right)^2 = O_p(NT).
\end{aligned}$$

Case (ii). Suppose that $k_1^0 \leq \hat{k} \leq k_2^0$. If $\hat{k} \in (k_1^0 + C_1, k_2^0]$, choosing $k \in [k_1^0, k_1^0 + C_1]$, we have

$$\begin{aligned}
U_{31}^{H_1} 1_{\{k > \hat{k}\}} &= 0, \quad U_{32}^{H_1} 1_{\{k > \hat{k}\}} = 0, \quad U_5^{H_1} = 0, \quad U_{10}^{H_1} = 0, \quad U_{11}^{H_1} = 0, \\
U_4^{H_1} &= \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=k_1^0+1}^k x'_{it} \delta_i^0 = O_p\left(\sqrt{\frac{N}{T}}\right),
\end{aligned}$$

since $k < \hat{k}$, and

$$U_{21}^{H_1} = \sqrt{\frac{T}{N}} \sum_{i \in G_1} \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \frac{1}{T} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 = O_p(\sqrt{NT}).$$

Thus, we have

$$\begin{aligned}
\sup_{k \in [1, T-1]} US_{NT}(k, \hat{k}) &\geq \sup_{k \in [k_1^0, k_1^0 + C_1]} US_{NT}(k, \hat{k}) \\
&= \sup_{k \in [k_1^0, k_1^0 + C_1]} \left(U_{21}^{H_1} + O_p(1) + O_p\left(\sqrt{\frac{N}{T}}\right) \right)^2 = O_p(NT).
\end{aligned}$$

If $\hat{k} \in [k_1^0, k_1^0 + C_1]$, since $(\hat{k} - k_1^0)/T = O_p(1/T)$,

$$U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} = O_p\left(\sqrt{\frac{N}{T}}\right), \quad U_{31}^{H_1} 1_{\{\hat{k} > k_1^0\}} = O_p\left(\sqrt{\frac{N}{T}}\right), \quad U_4^{H_1} = O_p\left(\sqrt{\frac{N}{T}}\right), \quad U_5^{H_1} = O_p\left(\sqrt{\frac{N}{T}}\right).$$

Choosing $k = k_2^0$, we have $U_{11}^{H_1} = 0$, and

$$\begin{aligned}
& U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} - U_{10}^{H_1} \\
&= \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^{k_2^0} x'_{it} \left(\sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \delta_i^0 - \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^{k_2^0} x'_{it} \delta_i^0 \\
&= -\sqrt{\frac{T}{N}} \sum_{i \in G_2} \frac{1}{T} \sum_{t=\hat{k}+1}^{k_2^0} x'_{it} \left(\frac{1}{T} \sum_{t=\hat{k}+1}^T x_{it} x'_{it} \right)^{-1} \frac{1}{T} \sum_{t=k_2^0+1}^T x_{it} x'_{it} \delta_i^0 \\
&= O_p(\sqrt{NT}).
\end{aligned}$$

Thus, we have

$$\sup_{k \in [1, T-1]} US_{NT}(k, \hat{k}) \geq US_{NT}(k_2^0, \hat{k}) = \left(U_{32}^{H_1} - U_{10}^{H_1} + O_p(1) + O_p\left(\sqrt{\frac{N}{T}}\right) \right)^2 = O_p(NT).$$

Case (iii). Suppose that $k_1^0 < k_2^0 < \hat{k}$, then we have

$$U_{32}^{H_1} 1_{\{\hat{k} < k_2^0\}} = 0, U_{10}^{H_1} = 0, U_{11}^{H_1} = 0.$$

Choosing $k \in (k_1^0, k_1^0 + C_1]$, we can see that $U_{31}^{H_1} 1_{\{k > \hat{k}\}} = 0$, $U_5^{H_1} = 0$, $U_4^{H_1} = O_p(\sqrt{N/T})$, and

$$\begin{aligned}
U_{21}^{H_1} 1_{\{\hat{k} > k_1^0\}} &= \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^k x'_{it} \left(\sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 \\
&= \sqrt{\frac{T}{N}} \sum_{i=1}^N \frac{1}{T} \sum_{t=1}^k x'_{it} \left(\frac{1}{T} \sum_{t=1}^{\hat{k}} x_{it} x'_{it} \right)^{-1} \frac{1}{T} \sum_{t=k_1^0+1}^{\hat{k}} x_{it} x'_{it} \delta_i^0 \\
&= O_p(\sqrt{NT}).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\sup_{k \in [1, T-1]} US_{NT}(k, \hat{k}) &\geq \sup_{k \in (k_1^0, k_1^0 + C_1]} US_{NT}(k, \hat{k}) \\
&= \sup_{k \in (k_1^0, k_1^0 + C_1]} \left(U_{21}^{H_1} + O_p(1) + O_p\left(\sqrt{\frac{N}{T}}\right) \right)^2 = O_p(NT).
\end{aligned}$$

The proof of Proposition 2 is complete. \blacksquare

Proof of Proposition 3. From Proposition 1, under the alternative H_{1A} , the estimated

common break \hat{k} takes a value in $[k_1^0 - C_1, k_2^0 + C_2]$ with probability approaching one, for arbitrary positive constants C_1, C_2 . Thus, we investigate the limiting properties of the normalization factor in three cases that $k_1^0 - C_1 \leq \hat{k} < k_1^0$, $k_1^0 \leq \hat{k} \leq k_2^0$, and $k_2^0 < \hat{k} \leq k_2^0 + C_2$.

Case (i). Suppose that $k_1^0 - C_1 \leq \hat{k} < k_1^0$, we have,

$$\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) \leq V_{NT}(k_1, \hat{k}, k_2^0), \text{ for } k_1 \in \Omega(\epsilon).$$

To show that the minimum value of $V_{NT}(k_1, \hat{k}, k_2)$ is stochastically bounded, it is sufficient to show that for any $k_1 \in \Omega(\epsilon)$,

$$V_{NT}(k_1, \hat{k}, k_2^0) = O_p(1).$$

In this case, the model is estimated by regressing Y_i on $[X_i, X_{1i}(k_1, \hat{k}), X_{2i}(\hat{k}, k_2^0), X_{3i}(k_2^0)]$, which is expressed as

$$\begin{aligned} Y_i &= [X_i, X_{1i}(k_1, \hat{k}), X_{2i}(\hat{k}, k_2^0), X_{3i}(k_2^0)] \begin{bmatrix} \beta_i \\ \delta_{1i} \\ \delta_{2i} \\ \delta_{3i} \end{bmatrix} + u_i \\ &= \tilde{X}_i(k_1, \hat{k}, k_2^0) b_{1i} + u_i, \end{aligned} \tag{B.33}$$

while the true model with distinct common breaks is defined by

$$\begin{aligned} Y_i &= [X_i, X_{1i}(\hat{k}, k_1^0), X_{2i}(k_1^0, k_2^0), X_{3i}(k_2^0)] b_{1i}^0 + u_i \\ &= \tilde{X}_i(\hat{k}, k_1^0, k_2^0) b_{1i}^0 + u_i \\ b_{1i}^0 &= \begin{cases} [\beta_i^{0'}, 0, \delta_i^{0'}, \delta_i^{0'}]' & \text{if } i \in G_1, \\ [\beta_i^{0'}, 0, 0, \delta_i^{0'}]' & \text{if } i \in G_2. \end{cases} \end{aligned} \tag{B.34}$$

Replacing Y_i in (B.33) by (B.34), the residuals can be written by, for individuals in group 1,

$$\begin{aligned} \tilde{u}_i &= \tilde{X}_i(\hat{k}, k_1^0, k_2^0) b_{1i}^0 + u_i - \tilde{X}_i(k_1, \hat{k}, k_2^0) \tilde{b}_{1i}(\hat{k}) \\ &= u_i - \tilde{X}_i(k_1, \hat{k}, k_2^0) [\tilde{b}_{1i}(\hat{k}) - b_{1i}^0] + [\tilde{X}_i(\hat{k}, k_1^0, k_2^0) - \tilde{X}_i(k_1, \hat{k}, k_2^0)] b_{1i}^0 \\ &= u_i - \tilde{X}_i(k_1, \hat{k}, k_2^0) \begin{bmatrix} \tilde{\beta}_i - \beta_i^0 \\ \tilde{\delta}_{1i} \\ \tilde{\delta}_{2i} - \delta_i^0 \\ \tilde{\delta}_{3i} - \delta_i^0 \end{bmatrix} + [0, X_{1i}(\hat{k}, k_1^0) - X_{1i}(k_1, \hat{k}), X_{2i}(k_1^0, k_2^0) - X_{2i}(\hat{k}, k_2^0), 0] \begin{bmatrix} \beta_i^0 \\ 0 \\ \delta_i^0 \\ \delta_i^0 \end{bmatrix} \\ &= u_i - X_i(\tilde{\beta}_i - \beta_i^0) - X_{1i}(k_1, \hat{k}) \tilde{\delta}_{1i} - X_{2i}(\hat{k}, k_2^0) (\tilde{\delta}_{2i} - \delta_i^0) - X_{3i}(k_2^0) (\tilde{\delta}_{3i} - \delta_i^0) \\ &\quad + [X_{2i}(k_1^0, k_2^0) - X_{2i}(\hat{k}, k_2^0)] \delta_i^0. \end{aligned} \tag{B.35}$$

For individuals in group 2, we have

$$\begin{aligned}\tilde{u}_i &= u_i - \tilde{X}_i(k_1, \hat{k}, k_2^0) \begin{bmatrix} \tilde{\beta}_i - \beta_i^0 \\ \tilde{\delta}_{1i} \\ \tilde{\delta}_{2i} \\ \tilde{\delta}_{3i} - \delta_i^0 \end{bmatrix} + [0, X_{1i}(\hat{k}, k_1^0) - X_{1i}(k_1, \hat{k}), X_{2i}(k_1^0, k_2^0) - X_{2i}(\hat{k}, k_2^0), 0] \begin{bmatrix} \beta_i^0 \\ 0 \\ 0 \\ \delta_i^0 \end{bmatrix} \\ &= u_i - X_i(\tilde{\beta}_i - \beta_i^0) - X_{1i}(k_1, \hat{k})\tilde{\delta}_{1i} - X_{2i}(\hat{k}, k_2^0)\tilde{\delta}_{2i} - X_{3i}(k_2^0)(\tilde{\delta}_{3i} - \delta_i^0).\end{aligned}\tag{B.36}$$

By the definition of the denominator, $V_{NT}(k_1, \hat{k}, k_2^0)$ can be decomposed into four parts $V_1^{H_1}$, $V_2^{H_1}$, $V_3^{H_1}$, and $V_4^{H_1}$, defined by

$$\begin{aligned}V_1^{H_1} &= \frac{1}{T} \sum_{s=1}^{k_1} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=1}^s \tilde{u}_{it} \right)^2, \quad V_2^{H_1} = \frac{1}{T} \sum_{s=k_1+1}^{\hat{k}} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^{\hat{k}} \tilde{u}_{it} \right)^2, \\ V_3^{H_1} &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2^0} \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s \tilde{u}_{it} \right)^2, \quad V_4^{H_1} = \frac{1}{T} \sum_{s=k_2^0+1}^T \left(\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=s}^T \tilde{u}_{it} \right)^2.\end{aligned}$$

From (B.35) and (B.36), for $t \leq \hat{k}$, the residuals \tilde{u}_{it} are calculated on the basis of subsamples $\{x_{i1}, \dots, x_{ik_1}\}$, and $\{x_{i(k_1+1)}, \dots, x_{i\hat{k}}\}$, which are the same as those in (A.22) under the null hypothesis. Using the asymptotic distribution of (A.23)–(A.27) and $k_1^0 - \hat{k} = O_p(1)$, we can derive the limiting distributions of the terms $V_1^{H_1}$ and $V_2^{H_1}$ as follows:

$$V_1^{H_1} + V_2^{H_1} \Rightarrow \sigma^2 \int_0^{\tau_1} \left(W(r) - \frac{r}{\tau_1} W(\tau_1) \right)^2 dr + \sigma^2 \int_{\tau_1}^{\tau_1^0} \left[W(\tau_1^0) - W(r) - \frac{\tau_1^0 - r}{\tau_1^0 - \tau_1} (W(\tau_1^0) - W(\tau_1)) \right]^2.$$

We next consider the third term, which can be rewritten as

$$\begin{aligned}V_3^{H_1} &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2^0} \left[\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s x'_{it}(\tilde{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=\hat{k}+1}^s x'_{it}(\tilde{\delta}_{2i} - \delta_i^0) \right. \\ &\quad \left. - \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \left(\sum_{t=\hat{k}+1}^s x'_{it} \delta_i^0 1_{\{s \leq k_1^0\}} + \sum_{t=\hat{k}+1}^{k_1^0} x'_{it} \delta_i^0 1_{\{s > k_1^0\}} \right) - \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^s x'_{it} \tilde{\delta}_{2i} \right]^2 \tag{B.37} \\ &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2^0} \left[\frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s u_{it} - \frac{1}{\sqrt{NT}} \sum_{i=1}^N \sum_{t=\hat{k}+1}^s x'_{it}(\tilde{\beta}_i - \beta_i^0) - \frac{1}{\sqrt{NT}} \sum_{i \in G_1} \sum_{t=\hat{k}+1}^s x'_{it}(\tilde{\delta}_{2i} - \delta_i^0) \right. \\ &\quad \left. - O_p \left(\sqrt{\frac{N}{T}} \right) - \frac{1}{\sqrt{NT}} \sum_{i \in G_2} \sum_{t=\hat{k}+1}^s x'_{it} \tilde{\delta}_{2i} \right]^2 \\ &= \frac{1}{T} \sum_{s=\hat{k}+1}^{k_2^0} \left(V_{31}^{H_1} - V_{32}^{H_1} - V_{33}^{H_1} - O_p \left(\sqrt{\frac{N}{T}} \right) - V_{34}^{H_1} \right)^2.\end{aligned}$$

Since $k_1^0 - \hat{k} = O_p(1)$, the terms in parentheses in (B.37) are $o_p(1)$ and will vanish as $N, T \rightarrow \infty$. Similar to V_{31} and V_{32} , we have

$$V_{31}^{H_1} \Rightarrow \sigma(W(r) - W(\tau_1^0)), \quad (\text{B.38})$$

$$V_{32}^{H_1} \Rightarrow \sigma(r - \tau_1^0) \frac{W(\tau_1)}{\tau_1}. \quad (\text{B.39})$$

The coefficient estimator $\tilde{\delta}_{2i}$ is calculated by, for $i \in G_1$,

$$\begin{aligned} \tilde{\delta}_{2i} &= \left(\sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} y_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} y_{it} \\ &= \left(\sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \left[\sum_{t=\hat{k}+1}^{k_1^0} x_{it} (x'_{it} \beta_i^0 + u_{it}) + \sum_{t=k_1^0+1}^{k_2^0} x_{it} (x'_{it} \beta_i^0 + x'_{it} \delta_i^0 + u_{it}) \right] \\ &\quad - \left[\beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &= \delta_i^0 + \left(\sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} - O_p\left(\frac{1}{T}\right), \end{aligned}$$

for $i \in G_2$,

$$\begin{aligned} \tilde{\delta}_{2i} &= \left(\sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} (x'_{it} \beta_i^0 + u_{it}) - \left[\beta_i^0 + \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} \right] \\ &= \left(\sum_{t=\hat{k}+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \sum_{t=\hat{k}+1}^{k_2^0} x_{it} u_{it} - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it}. \end{aligned}$$

Then, we have

$$\begin{aligned} V_{33}^{H_1} + V_{34}^{H_1} &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{1}{T} \sum_{t=k_1^0+1}^s x'_{it} + O_p\left(\frac{1}{T}\right) \right) \sqrt{T} \left[\left(\sum_{t=k_1^0+1}^{k_2^0} x_{it} x'_{it} \right)^{-1} \sum_{t=k_1^0+1}^{k_2^0} x_{it} u_{it} \right. \\ &\quad \left. - \left(\sum_{t=1}^{k_1} x_{it} x'_{it} \right)^{-1} \sum_{t=1}^{k_1} x_{it} u_{it} + O_p\left(\frac{1}{T}\right) + O_p\left(\frac{1}{\sqrt{T}}\right) + O_p\left(\frac{1}{T}\right) \right] \\ &\Rightarrow \sigma(r - \tau_1^0) \left(\frac{W(\tau_2^0) - W(\tau_1^0)}{\tau_2^0 - \tau_1^0} - \frac{W(\tau_1)}{\tau_1} \right). \end{aligned}$$

Thus, we can find that the limiting distribution of $V_3^{H_1}$ is

$$\sigma^2 \int_{\tau_1^0}^{\tau_2^0} \left[W(r) - W(\tau_1^0) - \frac{r - \tau_1^0}{\tau_2^0 - \tau_1^0} (W(\tau_2^0) - W(\tau_1^0)) \right]^2 dr.$$

Since the coefficient estimator $\hat{\delta}_{3i}$ remains the same in groups 1 and 2, we have

$$\begin{aligned} V_4^{H_1} &= \frac{1}{T} \sum_{s=k_2^0+1}^T \left\{ \frac{1}{\sqrt{NT}} \sum_{i=1}^N \left[\sum_{t=s}^T u_{it} - \sum_{t=s}^T x'_{it} (\hat{\beta}_{1i} - \beta_i^0) - \sum_{t=s}^T x'_{it} (\hat{\delta}_{3i} - \delta_i^0) \right] \right\}^2 \\ &\Rightarrow \sigma^2 \int_{\tau_2^0}^1 \left[W(1) - W(r) - \frac{1-r}{1-\tau_2^0} (W(1) - W(\tau_2^0)) \right]^2 dr. \end{aligned}$$

Thus, we can say that

$$\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) \leq V_{NT}(k_1, \hat{k}, k_2^0) = V_1^{H_1} + V_2^{H_1} + V_3^{H_1} + V_4^{H_1} = O_p(1).$$

The proof of Proposition 2(i) is complete.

Case (ii) Suppose that $k_1^0 \leq \hat{k} \leq k_2^0$. In this case, we have

$$\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) \leq V_{NT}(k_1^0, \hat{k}, k_2^0).$$

We can easily find that the term $V_{NT}(k_1^0, \hat{k}, k_2^0)$ estimated using true break points will have a finite limiting distribution.

Case (iii) Suppose that $k_2^0 < \hat{k} \leq k_2^0 + C_2$. In this case, from (B.27), we have $\hat{k} - k_2^0 = O_p(1)$.

Similar to the proof of case (i), we can show that

$$\inf_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}(k_1, \hat{k}, k_2) \leq V_{NT}(k_1^0, \hat{k}, k_2) = O_p(1), \text{ for any } k_2 \in \Omega(\epsilon).$$

Thus, we complete the proof of Proposition 3. ■

Proof of Theorem 2. From Proposition 1, we show that $P(\hat{k} \in [k_1^0 - C_1, k_2^0 + C_2]) \rightarrow 1$.

Furthermore, for any $\hat{k} \in [k_1^0 - C_1, k_2^0 + C_2]$,

$$\begin{aligned} \sup_{k \in \Omega(\epsilon)} US_{NT}(k, \hat{k}) &= O_p(NT), \\ \sup_{(k_1, k_2) \in \Omega(\epsilon)} V_{NT}^{-1}(k_1, \hat{k}, k_2) &= O_p(1) \text{ (or } \infty), \end{aligned}$$

from Propositions 2 and 3. Thus, the proof of Theorem 2 is complete. ■

Table 1: Critical values

c_{τ^0}	10%	5%	1%
$c_{0.1}$	43.425	56.822	92.840
$c_{0.2}$	43.912	57.249	92.341
$c_{0.3}$	45.501	57.962	93.689
$c_{0.4}$	45.427	57.997	90.335
$c_{0.5}$	45.540	57.842	85.984
$c_{0.6}$	45.250	57.276	90.397
$c_{0.7}$	46.489	59.175	93.728
$c_{0.8}$	45.201	59.248	94.886
$c_{0.9}$	43.515	57.203	92.908

Table 2: Size of the test DGP.1

T	N	10%	5%	1%
(a) $\rho = 0$				
20	10	0.145	0.089	0.034
	50	0.136	0.074	0.026
	100	0.098	0.053	0.015
50	10	0.086	0.048	0.011
	50	0.076	0.036	0.009
	100	0.063	0.027	0.006
100	10	0.073	0.032	0.005
	50	0.072	0.034	0.006
	100	0.064	0.033	0.004
200	10	0.083	0.037	0.008
	50	0.075	0.032	0.006
	100	0.086	0.041	0.008
(b) $\rho = 0.4$				
20	10	0.226	0.146	0.060
	50	0.234	0.153	0.069
	100	0.231	0.143	0.058
50	10	0.134	0.068	0.022
	50	0.151	0.084	0.028
	100	0.145	0.084	0.024
100	10	0.113	0.063	0.017
	50	0.105	0.058	0.016
	100	0.101	0.055	0.016
200	10	0.107	0.048	0.013
	50	0.091	0.043	0.009
	100	0.091	0.050	0.013

Table 3: Power of the test DGP.2 (under H_{1A})

T	N	10%	5%	1%
(a) $\rho = 0$				
20	10	0.149	0.088	0.021
	50	0.586	0.455	0.222
	100	0.846	0.741	0.474
50	10	0.281	0.174	0.046
	50	0.918	0.847	0.614
	100	0.993	0.982	0.911
100	10	0.561	0.425	0.190
	50	0.996	0.980	0.916
	100	1.000	1.000	0.992
(b) $\rho = 0.4$				
20	10	0.304	0.216	0.100
	50	0.803	0.722	0.519
	100	0.965	0.929	0.789
50	10	0.390	0.276	0.121
	50	0.950	0.901	0.742
	100	0.997	0.991	0.946
100	10	0.611	0.491	0.258
	50	0.994	0.985	0.932
	100	1.000	1.000	0.994

¹ $k_1^0 = [T/4]$, $k_2^0 = [3T/4]$,
 $N_1 : N_2 = 5 : 5$.

Table 4: Power of the test DGP.2 (under H_{1A})

ρ	δ_{1i}, δ_{2i}	10%	5%	1%
0	U(0,0.1)	0.130	0.064	0.012
	U(0.1,0.2)	0.538	0.401	0.167
	U(0.2,0.3)	0.918	0.850	0.611
	U(0.3,0.4)	0.995	0.984	0.919
	U(0.4,0.5)	1.000	0.999	0.986
	U(0.5,0.6)	1.000	1.000	0.998
	U(0.6,0.7)	1.000	1.000	1.000
	U(0.7,0.8)	1.000	1.000	1.000
	U(0.8,0.9)	1.000	1.000	1.000
	U(0.9,1.0)	1.000	1.000	1.000
	U(1.4,1.5)	1.000	1.000	1.000
0.4	U(0,0.1)	0.206	0.138	0.037
	U(0.1,0.2)	0.652	0.540	0.292
	U(0.2,0.3)	0.955	0.913	0.750
	U(0.3,0.4)	0.997	0.992	0.949
	U(0.4,0.5)	1.000	1.000	0.993
	U(0.5,0.6)	1.000	1.000	1.000
	U(0.6,0.7)	1.000	1.000	1.000
	U(0.7,0.8)	1.000	1.000	1.000
	U(0.8,0.9)	1.000	1.000	1.000
	U(0.9,1.0)	1.000	1.000	1.000
	U(1.4,1.5)	1.000	1.000	1.000

¹ $T = 50, N = 50$.

² $k_1^0 = [T/4], k_2^0 = [3T/4], N_1 : N_2 = 5 : 5$.

Table 5: Power of the test DGP.2 (under H_{1A})

k_1^0	k_2^0	10%	5%	1%
[0.2T]	[0.25T]	0.120	0.066	0.018
	[0.3T]	0.325	0.225	0.070
	[0.4T]	0.801	0.695	0.465
	[0.5T]	0.941	0.891	0.734
	[0.6T]	0.955	0.918	0.764
	[0.7T]	0.925	0.875	0.692
	[0.8T]	0.859	0.771	0.558

¹ $N = T = 50$, $\rho = 0.4$, $N_1 : N_2 = 5 : 5$.

Table 6: Power of the test DGP.2 (under H_{1A})

$N_1 : N_2$	10%	5%	1%
2:N-2	0.168	0.105	0.037
1:9	0.262	0.187	0.073
2:8	0.546	0.447	0.241
3:7	0.811	0.719	0.500
4:6	0.938	0.888	0.737
5:5	0.978	0.950	0.840

¹ $N = T = 50, \rho = 0.4$.

² $k_1^0 = [0.3T], k_2^0 = [0.7T]$.

Table 7: Power of the test ($\rho = 0.4$ under H_{2A})

k_1^0	k_2^0	k_3^0	$N_1 : N_2 : N_3$	T	N	10%	5%	1%
[T/6]	[3T/6]	[4T/6]	3:3:4	50	10	0.296	0.204	0.080
					50	0.646	0.522	0.284
					100	0.740	0.642	0.445
[0.4T]	[0.5T]	[0.6T]	3:3:4	50	10	0.248	0.165	0.058
					50	0.485	0.381	0.211
					100	0.629	0.506	0.296
[0.2T]	[0.25T]	[0.5T]	3:3:4	50	10	0.342	0.239	0.109
					50	0.835	0.750	0.562
					100	0.964	0.923	0.809
[0.2T]	[0.3T]	[0.8T]	3:3:4	50	10	0.334	0.235	0.087
					50	0.704	0.591	0.363
					100	0.851	0.772	0.583
[0.2T]	[0.5T]	[0.8T]	1:4:5	50	10	0.334	0.224	0.087
					50	0.851	0.758	0.566
					100	0.925	0.866	0.725

Table 8: Power of the test (under H_{3A})

T	N	10%	5%	1%
(a) $\rho = 0$				
20	10	0.147	0.089	0.022
	50	0.548	0.412	0.191
	100	0.654	0.497	0.255
50	10	0.281	0.173	0.050
	50	0.688	0.515	0.247
	100	0.884	0.770	0.474
100	10	0.468	0.329	0.129
	50	0.908	0.815	0.541
	100	0.969	0.895	0.637
(b) $\rho = 0.4$				
20	10	0.344	0.243	0.109
	50	0.725	0.584	0.352
	100	0.842	0.740	0.493
50	10	0.300	0.204	0.076
	50	0.842	0.730	0.477
	100	0.903	0.810	0.538
100	10	0.517	0.392	0.183
	50	0.942	0.874	0.642
	100	0.879	0.761	0.438

Table 9: Power of the test DGP.3 (under H_{4A})

T	N	10%	5%	1%
(a) $\rho = 0$				
20	10	0.316	0.219	0.087
	50	0.701	0.600	0.374
	100	0.876	0.831	0.685
50	10	0.555	0.428	0.218
	50	0.964	0.939	0.807
	100	0.996	0.994	0.973
100	10	0.771	0.676	0.448
	50	0.999	0.995	0.981
	100	1.000	1.000	1.000
(b) $\rho = 0.4$				
20	10	0.552	0.447	0.267
	50	0.920	0.867	0.706
	100	0.991	0.975	0.914
50	10	0.684	0.571	0.351
	50	0.994	0.984	0.908
	100	0.999	0.999	0.987
100	10	0.815	0.728	0.531
	50	0.999	0.998	0.984
	100	1.000	1.000	1.000

¹ $k_1^0 = [T/4]$, $k_2^0 = [3T/4]$.

Table 10: Detection of common break dates during the period 2005M02 – 2011M12

Category	N	T	Statistic $S_{NT}(k, k_1, k_2)$	Estimated common break date
(a) 2005M02 – 2011M12				
Foreign Large Blend	58	83	270.2507***	2008M01
Foreign Small/Mid Blend	7	83	98.1177***	2008M02
Foreign Large Growth	39	83	299.1527***	2008M01
Foreign Small/Mid Growth	11	83	8.5283	2008M11
Large Blend	205	83	278.6359***	2008M01
Mid-Cap Blend	54	83	305.5441***	2008M01
Small Blend	76	83	240.9046***	2008M01
Large Growth	186	83	100.4435***	2008M02
Mid-Cap Growth	88	83	103.6767***	2008M02
Small Growth	92	83	110.178***	2008M02
(b) 2005M02 – 2008M05				
Foreign Large Blend	58	40	146.0774***	2007M12
Foreign Small/Mid Blend	7	40	146.2134***	2007M12
Foreign Large Growth	39	40	145.7293***	2007M12
Foreign Small/Mid Growth	11	40	116.7004***	2007M12
Large Blend	205	40	150.0121***	2007M12
Mid-Cap Blend	54	40	146.7684***	2007M12
Small Blend	76	40	137.2884***	2007M12
Large Growth	186	40	157.4825***	2007M12
Mid-Cap Growth	88	40	151.1206***	2007M12
Small Growth	92	40	136.7168***	2007M12
(c) 2008M06 – 2011M12				
Foreign Large Blend	58	43	52.1541*	2008M12
Foreign Small/Mid Blend	7	43	26.9158	2008M12
Foreign Large Growth	39	43	31.5376	2008M12
Foreign Small/Mid Growth	11	43	34.1773	2008M12
Large Blend	205	43	10.6933	2008M12
Mid-Cap Blend	54	43	9.4507	2008M12
Small Blend	76	43	13.8338	2008M12
Large Growth	186	43	16.8967	2008M12
Mid-Cap Growth	88	43	20.9836	2008M12
Small Growth	92	43	9.853	2008M12

¹ * reject at the 10% significance level.

² ** reject at the 5% significance level.

³ *** reject at the 1% significance level